

## Sitting still or going hunting: Which works better?

November 1 2012, by David Chandler



Detailed computer simulation shows how a particle of nutrients gets distributed by turbulent water. Image courtesy of courtesy of John R. Taylor and Roman Stocker

For the kinds of animals that are most familiar to us—ones that are big enough to see—it's a no-brainer: Is it better to sit around and wait for food to come to you, or to move around and find it? Larger animals that



opt to sit around aren't likely to last long.

But for bacteria out in the ocean, the question is a far more complicated one.

<u>Oceanographers</u> have long assumed that because turbulence distributes nutrients uniformly through the water, and because the ability of tiny organisms to move around is insignificant compared to this turbulence, there was no reason for such creatures to move at all. Sea-dwelling bacterial life, they believed, should consist just of static feeders.

That view has now been upended by research conducted by Roman Stocker, an associate professor in MIT's Department of Civil and Environmental Engineering, and John R. Taylor, a former MIT postdoc who is now a lecturer in applied mathematics and <u>theoretical physics</u> at Cambridge University. It turns out that swimmers and passive feeders each have some advantages—but also pay some costs—in food gathering, depending on how fast the swimmers swim and how strong the turbulence is.

The results, based on a <u>computer model</u> that for the first time considers nutrient competition by bacteria in a <u>turbulent flow</u>, have just been published in *Science*.

Until now, most studies of microorganism behavior have taken place in static environments, such as a <u>test tube</u> or Petri dish, without regard to <u>fluid flow</u>. "<u>Marine bacteria</u> have mostly been studied in isolation from the motion of the seawater they live in," Stocker says. These new <u>computer simulations</u>, which model both how water flows and how bacteria in that water behave, have now made it possible to study the foraging of bacteria in dynamic environments, similar to the turbulent waters they naturally inhabit.



"We're working at the interface between microbiology and <u>fluid</u> <u>dynamics</u>," Stocker says. The new work has produced the first study of how this turbulent environment affects the choice of bacterial feeding strategy.

While it's true that ocean currents and turbulence are orders of magnitude faster than anything that even the most energetic <u>microorganisms</u> can achieve, it turns out that these creatures' motions are nevertheless very important, Stocker and Taylor found. That's because the microenvironment a bacterium occupies—the microliters of water surrounding it and representing its foraging ground—is transported en masse in those flows, so even small rates of swimming can make a big difference in accessing food.

Although swimming toward a food source can confer some advantage in the amount of food consumed, it also carries a penalty: Swimming uses energy, requiring more food. "As you swim faster, you do better in terms of food uptake, but you also pay a price," Stocker says.

Because of that tradeoff, the researchers found, for any given intensity of turbulence there is an optimal swimming speed; anything faster than this produces diminishing returns.

"What we did was to quantify the cost-benefit trade-off," Stocker says. "We considered the pluses and minuses, and found that the swimming costs matter." In general, the results showed, the optimal swimming speed for marine bacteria is about 60 micrometers per second, which is "in very nice agreement with what's observed for the actual swimming speed of many marine bacteria," Stocker adds.

Turbulence stirs nutritious material through the water in much the same way as it stirs cream into a cup of coffee: First forming filaments, then mixing them away. "Something similar happens—on a tiny scale—when



bursts of organic matter enter gently moving water," Taylor says. "The swirls of organic matter are easily accessed by swimming bacteria, which surround and absorb it."

But the swimmers' advantage is short-lived. In turbulent <u>seawater</u>, after several seconds, nutrients are uniformly distributed—meaning there's as much available for the stationary microbes as there is for the swimmers.

Most of this organic matter is derived from phytoplankton, tiny organisms that capture sunlight and convert it into carbohydrates. In fact, phytoplankton in the ocean are responsible for about as much photosynthesis as all of Earth's terrestrial plant matter combined, so understanding their role in the global carbon cycle is crucially important for understanding everything from the food chains that sustain the world's fisheries to the complexities of global climate change.

As part of their analysis, Stocker and Taylor conducted detailed simulations of exactly how a patch of nutrient-rich material, derived from dying phytoplankton or other sources, disperses in turbulent water. What starts out as a compact blob "gets stretched out, stirred, and explodes into a tangled web of filaments," Stocker says.

That's when the swimming advantage is greatest: The filaments provide lots of nutrient-rich zones to swim toward. But then the window of opportunity closes. "Eventually it all gets homogenized," he says. "Once it's mixed, swimming makes no more sense".

How fast that happens depends on the strength of turbulence. Stocker and Taylor found that very high or very low turbulence provides the least advantage for the swimmers. But for a range in between, they had a substantial edge over their inert relatives.

The basic understanding of these microscopic processes could have large-



scale importance, Stocker says. As the Earth's climate warms, turbulence might become weaker in the ocean, which could affect which species dominate the competition for nutrients at the base of the food chain. Species composition, in turn, can dictate how much carbon is channeled to organisms higher in the food chain, shaping the productivity of a marine environment.

## Provided by Massachusetts Institute of Technology

Citation: Sitting still or going hunting: Which works better? (2012, November 1) retrieved 23 June 2024 from <u>https://phys.org/news/2012-11-sitting-still-or-going-hunting.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.