

Astronomers go infrared to map brightest galaxies in Universe

December 4 2012



A 3D projection of almost 300 galaxies in the census in the same part of the sky. The third dimension shows how many billions of years back in time we are seeing each galaxy, determined by observations from the Keck Observatory. At top are images from the Hubble Space Telescope of five galaxies in the census. Credit: ESA-C. Carreau

(Phys.org)—A group of astronomers from the University of Hawaii at Manoa, the U.S. Mainland, Canada, and Europe recently used the twin telescopes of the W. M. Keck Observatory on Mauna Kea, Hawaii, to conduct a census of the brightest, but until now unseen, galaxies in the distant Universe, bringing astronomers one step closer to understanding



how galaxies form and evolve.

These <u>galaxies</u> glow so brightly at <u>infrared wavelengths</u> that they would outshine our own Milky Way by hundreds, maybe thousands, of times. They are forming stars so quickly that between 100 and 500 new stars are born in each galaxy every year, and have been coined "starbursts" by astronomers.

While it's not clear what gives these galaxies their intense luminosity, it could be the result of a collision between two spiral-type galaxies, similar to the Milky Way and Andromeda Galaxies. Or they could be in a particularly gas-rich region of space, where galaxies form stars quickly due to constant bombardment from gas and dust.

Despite their brightness, these galaxies are nearly invisible at the wavelengths our eyes and most telescopes on Earth can see because they contain huge amounts of dust, which absorbs visible starlight. But they were detectable directly in the infrared from observations at the Herschel Space Observatory, said Dr. Caitlin Casey, a Hubble fellow at the UH Manoa Institute for Astronomy and the lead scientist behind the new results. "Herschel is an <u>infrared space telescope</u> sensitive to wavelengths not observable from within Earth's atmosphere," she said.





Images of six of the dusty galaxies as seen by Hubble Space Telescope with distances (in billions of light years) measured by the W. M. Keck Observatory. The background is an image of the sky in far-infrared from Herschel, where dusty galaxies like this are most easily detected. The Keck data (shown in blue below galaxy images), which spreads the light of each galaxy into a spectrum, allowed the astronomers to measure distances to each galaxy using "fingerprints" of atoms and molecules in each galaxy. These galaxies were previously unidentified due to the large amounts of dust absorbing the visible light. Credit: ESA-C. Carreau



"Detecting these bright <u>infrared galaxies</u> used to be difficult, and a handful was plenty; now with Herschel we are finding them by the thousands, enabling a census like this," said Göran Pilbratt, <u>European</u> <u>Space Agency</u> Herschel Project Scientist.

Once found, taking measurements of these galaxies at <u>visible</u> <u>wavelengths</u> required using the 10-meter Keck telescopes, the two largest <u>optical telescopes</u> in the world. Over the course of several nights the group was able to detect and measure distances to nearly 800 of these galaxies.

"For the first time, we have been able to measure distances, star formation rates, and temperatures for a brand new set of 767 previously unidentified galaxies," said Dr. Scott Chapman, a co-author on the studies. "The previous similar survey of distant infrared starbursts only covered 73 galaxies. This is a huge improvement."

"While some of the galaxies are nearby, most are very distant; we even found galaxies that are so far that their light has taken 12 billion years to travel here, so we are seeing them when the Universe was only a ninth of its current age," Casey said. "Now that we have a pretty good idea of how important this type of galaxy is in forming huge numbers of stars in the Universe, the next step is to figure out why and how they formed."

"It's hard to figure out how most galaxies formed based on information from only a small part of the Universe, just like it's hard to guess how big an elephant is if you only get a glimpse of its tail," Casey said. "Now that we have an accurate census of starbursting galaxies across a huge time period in the Universe's history, we can start to piece together how these galaxies grew and evolved."

Two papers detailing these results are published online today in the *Astrophysical Journal*.



More information: Casey, C.M., et al. A redshift survey of Herschel far-infrared selected starbursts and implications for obscured star formation, *The Astrophysical Journal*, 04 Dec 2012.

Casey, C.M., et al. A population of z>2 far-infrared Herschel-SPIRE selected starbursts, *The Astrophysical Journal*, 04 Dec 2012.

Provided by W. M. Keck Observatory

Citation: Astronomers go infrared to map brightest galaxies in Universe (2012, December 4) retrieved 2 May 2024 from <u>https://phys.org/news/2012-12-astronomers-infrared-brightest-galaxies-universe.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.