


GRAIL creates most accurate Moon gravity map (w/ video)

December 5 2012



These maps of the moon show the "Bouguer" gravity anomalies as measured by NASA's GRAIL mission. Bouguer gravity is what remains from the gravity field when the attraction of surface topography is removed, and therefore represents mass anomalies inside the moon due to either variations in crustal thickness or crust or mantle density. Red areas have stronger gravity, while blue areas have weaker gravity. Image credit: NASA/JPL-Caltech/CSM

(Phys.org)—Twin NASA probes orbiting Earth's moon have generated the highest resolution gravity field map of any celestial body.

The new map, created by the Gravity Recovery and Interior Laboratory (GRAIL) mission, is allowing scientists to learn about the moon's internal structure and composition in unprecedented detail. Data from

the two washing machine-sized spacecraft also will provide a better understanding of how Earth and other [rocky planets](#) in the solar system formed and evolved.


The gravity field map reveals an abundance of features never before seen in detail, such as tectonic structures, volcanic landforms, basin rings, crater central peaks and numerous simple, bowl-shaped craters. Data also show the moon's gravity field is unlike that of any [terrestrial planet](#) in our solar system.

These are the first scientific results from the prime phase of the mission, and they are published in three papers in the journal *Science*.

"What this map tells us is that more than any other [celestial body](#) we know of, the moon wears its gravity field on its sleeve," said GRAIL Principal Investigator Maria Zuber of the Massachusetts Institute of Technology in Cambridge. "When we see a notable change in the gravity field, we can sync up this change with [surface topography](#) features such as craters, rilles or mountains."

According to Zuber, the moon's gravity field preserves the record of impact bombardment that characterized all terrestrial planetary bodies and reveals evidence for fracturing of the interior extending to the deep crust and possibly the mantle. This impact record is preserved, and now precisely measured, on the moon.


The probes revealed the bulk density of the moon's highland crust is substantially lower than generally assumed. This low-bulk crustal density agrees well with data obtained during the final Apollo lunar missions in the early 1970s, indicating that local samples returned by astronauts are indicative of global processes.



These maps of the near and far side of the moon show gravity gradients as measured by NASA's GRAIL mission. Red and blue areas indicate stronger gradients due to underlying mass anomalies. Image credit: NASA/JPL-Caltech/CSM

"With our new crustal bulk density determination, we find that the average thickness of the moon's crust is between 21 and 27 miles (34 and 43 kilometers), which is about 6 to 12 miles (10 to 20 kilometers) thinner than previously thought," said Mark Wieczorek, GRAIL co-investigator at the Institut de Physique du Globe de Paris. "With this crustal thickness, the bulk composition of the moon is similar to that of Earth. This supports models where the moon is derived from Earth materials that were ejected during a giant impact event early in solar system history."

The map was created by the spacecraft transmitting radio signals to define precisely the distance between them as they orbit the moon in formation. As they fly over areas of greater and lesser gravity caused by visible features, such as mountains and craters, and masses hidden beneath the lunar surface, the distance between the two spacecraft will change slightly.




A 300-mile-long (500 kilometer-long) linear gravity anomaly on the far side of the moon has been revealed by gravity gradients measured by NASA's GRAIL mission. GRAIL data are shown on the left, with red and blue corresponding to stronger gravity gradients. Topography data from NASA's Lunar Reconnaissance Orbiter's Lunar Orbiter Laser Altimeter for the same region show a surface saturated with craters (red is higher terrain and blue is lower). The lack of any topographic signature over the gravity anomaly indicates that it is older than the craters, and thus is one of the oldest features on the moon. The units of the gravity gradients are Eötvös, and of the topography are kilometers. Image credit: NASA/JPL-Caltech/CSM

"We used gradients of the [gravity field](#) in order to highlight smaller and narrower structures than could be seen in previous datasets," said Jeff Andrews-Hanna, a GRAIL guest scientist with the Colorado School of Mines in Golden. "This data revealed a population of long, linear gravity anomalies, with lengths of hundreds of kilometers, crisscrossing the surface. These linear gravity anomalies indicate the presence of dikes, or long, thin, vertical bodies of solidified magma in the subsurface. The dikes are among the oldest features on the moon, and understanding

them will tell us about its early history."

While results from the primary science mission are just beginning to be released, the collection of gravity science by the lunar twins continues. GRAIL's extended mission science phase began Aug. 30 and will conclude Dec. 17. As the end of mission nears, the spacecraft will operate at lower orbital altitudes above the [moon](#).



A profile across one of the linear gravity anomalies found by NASA's GRAIL mission shows that it has higher gravity than the surroundings (the anomaly is the red line in graph at bottom left; its location is shown as a blue line in the center of the GRAIL gravity gradient map at top left). Models show that this gravity pattern indicates the presence of giant dikes beneath the surface of the moon (white line in graph at bottom left). A dike is a solidified magma filled crack that forms beneath the surface. Dikes on Earth are sometimes exposed by erosion so that they are visible on the surface, as seen at Ship Rock, New Mexico, pictured here at right. The lunar dikes identified by GRAIL are 50 times longer and 1,000 times wider than the dike seen here. Image credits, Left: NASA/JPL-Caltech/CSM, Right: Photo Copyright © Louis Maher

When launched in September 2011, the probes were named GRAIL A and B. They were renamed Ebb and Flow in January by elementary students in Bozeman, Mont., in a nationwide contest. Ebb and Flow were placed in a near-polar, near-circular orbit at an altitude of approximately 34 miles (55 kilometers) on Dec. 31, 2011, and Jan. 1, 2012, respectively.

NASA's Jet Propulsion Laboratory in Pasadena, Calif., manages the mission for NASA's Science Mission Directorate in Washington. GRAIL is part of the Discovery Program managed at NASA's Marshall Space Flight Center in Huntsville, Ala. Lockheed Martin Space Systems of Denver built the spacecraft.

Provided by NASA

Citation: GRAIL creates most accurate Moon gravity map (w/ video) (2012, December 5)
retrieved 1 May 2024 from

<https://phys.org/news/2012-12-grail-accurate-moon-gravity-video.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--