

Masers in stellar nurseries

December 24 2012



A false-color infrared image of a young star showing outflowing jets as green beams of shocked gas (actually two of the stars in this image have jets). A new study using the SMA has found that bright methanol masers, often seen in such star-forming regions and long thought to indicate only the youngest such stars, are also found around more mature stars. Credit: NASA-Spitzer Space Telescope

(Phys.org)—Astronomers have come to realize that the process of star formation, once thought to consist essentially of just the simple coalescence of material by gravity, occurs in a complex series of stages.

As the gas and dust in giant molecular clouds comes together into stars, dramatic outflowing jets of material develop around each, as do circumstellar disks (possibly pre-planetary in nature). Other features are present as well: Astronomers in the 1960s were amazed to discover that these star-forming regions sometimes produce natural masers (masers are the bright, radio wavelength analogs of lasers). Clouds of water vapor or methanol vapor in regions of active star formation generate some of the most spectacular masers.

Although associated with the complex activity of star formation, the role of masers in the building of a [new star](#) is thought to be minor (although it is not understood). However masers, because they are so bright, provide valuable diagnostic probes of the regions where star formation is underway. Exactly what they reveal is less clear, but many astronomers have thought that methanol masers can signal the very earliest stages of star formation, perhaps less than about ten thousand years old. One of the key questions masers can possibly help resolve is how stars more massive than the Sun form. Understanding the birth of such [massive stars](#) is essential not only in its own right, but also because these stars end up as [supernovae](#) which enrich the cosmos with elements essential to life. The birth of massive stars is, however, notoriously tricky to understand because their larger masses prompt the young star to mature very quickly, in less than about one hundred thousand years and much faster than lower-mass stars. As a result, many growth stages are blurred together. Masers are thought to offer a way to probe these earliest times of star formation.

SAO astronomers Claudia Cyganowski and Qizhou Zhang, with five colleagues, used the Submillimeter Array (SMA) to study regions of massive star formation identified in infrared images as having outflows typical of massive young stars. The SMA was able to identify all of the protostellar cores from their millimeter dust emission. They found one such protocluster of young stars that also contained a variety of types of

methanol masers, enabling a comparative study of masers and star-formation activity. Writing in the latest issue of the *Astrophysical Journal Letters*, the scientists report finding that, contrary to the conventional wisdom, methanol masers thought to be associated with very young stages of [star formation](#) are found occurring with more evolved embryos. The new results show for the first time that the mechanisms at work to make these methanol masers, shocks for example, are found in a much wider range of situations than previously suspected. The new results are not atypical of progress in astronomy.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Masers in stellar nurseries (2012, December 24) retrieved 27 April 2024 from <https://phys.org/news/2012-12-masers-stellar-nurseries.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.