

New species of Trilophodont gomphotheres found from the quaternary of China

December 17 2012



Fig.1 The skull and mandible of *Sinomastodon jiangnanensis* sp. nov. (IVPP V 18221), lateral view, scale equals 10 cm. Credit: WANG Yuan

According to an article published online December 2012 in the journal of *Chinese Science Bulletin* (Vol.57, No.36), Dr. JIN Changzhu, Institute of Vertebrate Paleontology and Paleoanthropology (IVPP), Chinese Academy of Sciences, and his team described a new species of

trilophodont gomphotheres, *Sinomastodon jiangnanensis*, from the Early Pleistocene Renzidong Cave deposits in Anhui Province, Eastern China. The specimen, an almost complete skull with mandible and dentition, is the first discovery of a Quaternary *Sinomastodon* skull from China and has significant implications for discussing the evolution, dispersion and related paleoecological variation of *Sinomastodon* during the Quaternary.

Gomphotheres are considered as the main clade of the [mastodons](#) and was one of the most successful groups of Proboscidea due to its wide geographic distribution across all continents except Antarctic and Australia, from the Early Miocene to the Late [Pleistocene](#). Early members were characterized by long jaws that progressively evolved into the short-jawed types found in both the Old and New World.

As the only brevirostrine (short and broad-jawed) trilophodont gomphothere known from the Old World, *Sinomastodon* was mainly indigenous to China from the Early Pliocene to the Pleistocene, but only a few single Pleistocene teeth were previously found in China.




Fig.2 The molars and fragmentary upper incisor of *Sinomastodon jiangnanensis* sp. nov., scale equals 5 cm. Credit: WANG Yuan

The Renzidong Paleolithic site discovered in 1998 is located near the south bank of the Yangtze River in Fanchang County, Anhui Province, Eastern China. This location lies in the transitional zone between the Palaearctic and Oriental zoogeographical regions. Systematic excavations of the cave have yielded numerous mammalian remains in association with stone and bone artifacts. Preliminary paleomagnetic dating shows that the age of the strata yielding *Sinomastodon* in Renzidong Cave is 2.14–2.15 million years ago.

With a brevirostrine, elephant-like skull, no lower tusks, and simple bunodont and trilophodont intermediate molars, the new species is morphologically distinct from other gomphotheres and should be assigned to the genus *Sinomastodon*.

Comparison studies indicate that the new species is more progressive than *S. hanjiangensis* and the Pliocene type species *S. intermedius* in its skull and mandible morphology, but is evidently more primitive than the Pleistocene *S. yangziensis* in its molar morphology.

"The faunal analysis of the Renzidong fauna indicates a warm semi-arid and semi-humid forest-grassland environment. Obviously, this fauna has characteristics of a transitional zone with more typical northern types", said WANG Yuan, first author of the study at the IVPP, "The large scale invasion of northern forms moving to the south implies a distinct cooling event at the beginning of the Quaternary, which resulted in a southward migration of *Sinomastodon* to the Jiangnan area and the speciation of *S. jiangnanensis*."

This work was supported by the [Chinese Academy of Sciences](#), the National Natural Science Foundation of China, the [China Geological Survey](#), and the Key Laboratory of Evolutionary Systematics of Vertebrates, CAS.

Provided by Institute of Vertebrate Paleontology and Paleoanthropology

Citation: New species of Trilophodont gomphotheres found from the quaternary of China (2012, December 17) retrieved 29 June 2024 from <https://phys.org/news/2012-12-species-trilophodont-gomphotheres-quaternary-china.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.