

Storms in the machine

December 18 2012



Moving west-northwest, Typhoon Imbudo approaches the southern coast of China on July 23, 2003. This Moderate Resolution Imaging Spectroradiometer (MODIS) image from the morning of July 23, shows the storm stretched out over the South China Sea. At that time, sustained winds were about 90 knots, or 104 mph. Credit: Jacques Descloitres, MODIS Rapid Response Team at NASA GSFC

NASA researchers are constructing a statistical typhoon model that can quickly generate millions of synthetic typhoons from birth through termination. Their work could provide new details about the climate of Earth, and what the future may hold for life on our planet.


The year is 1975, and a dark cloud is looming off the coast of Taiwan. With little warning, the storm comes crashing ashore with winds of over 150 miles per hour. The '[super typhoon](#)' tears its way across the island - but the real horror comes when it makes its way onto [Mainland China](#). Although weakened by its journey over the [mountains](#) of Taiwan, typhoon [Nina](#) lashes inland China with [torrential rain](#). This results in the catastrophic failure of the Banquiao Dam. Floods sweep over the land, claiming over 171,000 lives and causing billions of dollars in damage.

In 2011 alone, by some estimates the cost of "Typhoon Season" in the Pacific was over 4 billion US dollars and total deaths passed 1700. Many of the most populous cities in the world now line the coastal regions of the Asian mainland, Japan, the Philippines and other Pacific island states. In China alone, some 250 million people live in areas threatened by such storms. Every year, the lives of millions upon millions of people are in harms way.

Asian Brew

When a typhoon begins brewing in the Pacific, numerical weather modeling is used to forecast how the storm will evolve in the following days. These storms can develop with ferocious speed and quickly sweep toward populated areas.

Unfortunately, a few days warning is not enough to prevent the death and destruction that follows.



Landfall rates (landfalls per 100km of segmented coastline per year) as functions of distance along five sections of coast: mainland Asia, Japan, Philippines, Indonesia/Malaysia, and Kamchatka. Colors indicate ENSO phase; red is ENSO = +2 (strong El Niño) and blue ENSO = -2 (strong La Niña). Credit: NASA/GISS/Yonekura & Hall

Local governments, disaster management groups and the insurance industry would benefit from more time to plan and prepare. In fact, it takes these institutions years to develop effective methods for dealing with massive typhoons. They need to be able to answer questions like: what are the odds that a given coastal city will suffer a major typhoon strike in the next year? Or the next five years? How are these odds affected by our changing [climate](#)? How can past typhoon observations, trends, and fluctuations in [climate](#) be used to inform current and future policies in coastal regions? A new computer model may have some answers.

Typhoons in the Machine


Two researchers from NASA and Columbia University, Emmi Yonekura and Tim Hall, are constructing a statistical typhoon model that can quickly generate millions of synthetic typhoons from birth through termination, and whose statistical properties match those of the hundreds of historical documented typhoons. They then use the synthetic set to estimate local Asian landfall rates with a precision much higher than could be obtained directly from historical landfalls. In its current stage the model simulates typhoon formation and propagation and their dependencies on a natural climate pattern known as "El Niño/Southern Oscillation" (ENSO).



This spectacular, oblique view shows the eye of typhoon Odessa, southeast of Japan. An astronaut's perspective from low Earth orbit provides details of large storm systems that cannot be resolved from other meteorological satellites.

Credit: NASA/STS51I-37-0083

ENSO - called El Niño or La Niña depending on phase - is the [Earth's](#) dominant year-to-year climate variability. The ENSO cycle is irregular, repeating every 2 to 7 years. Put simply, ENSO is the east-west sloshing of warm tropical Pacific surface waters. ENSO not only affects Pacific-sector weather, but also has global influence on temperature, wind patterns, and rainfall.



Examples of simulated typhoon tracks in the northwest Pacific. The colored

segments show coastal regions used to compute landfall rates. Credit: NASA/GISS/Yonekura & Hall

Using carefully optimized statistical analysis of past typhoon behavior, Yonekura and Hall are able to quantify how the number of typhoons forming each year in the north Pacific is dependent on ENSO. Additionally, they have been able to gather data on where in the north Pacific the storms form and how, on average, the storms propagate. The researchers use these storms' dependencies on ENSO in their model to generate tens of thousands of typhoons in each of several fixed ENSO states spanning the observed range of ENSO. This allowed them to compute landfall rates of typhoons at high spatial resolution along Asian coastlines as a function of ENSO phase.

Predicting the future

The overall annual number of typhoons changes only modestly with ENSO, but there are significant changes in the shape of typhoon tracks and the dominant regions of typhoon formation. El Niño typhoons propagate further westward before re-curving northeast, a feature that increases their odds of making landfall. However, El Niño typhoons also form further eastward, away from the coast, which reduces their landfall odds. The combined effect is an overall reduction in landfall during El Niño and an increase during La Niña (although there is considerable variation along the coast). Southern China and Southeast Asia experience the largest increase of landfall in La Niña, while the Philippines show no overall ENSO sensitivity and even regions of opposite sensitivity.

As the Pacific nations brace for the 2013 typhoon season, meteorologists will ply their skills to forecast the evolution over days of particular storms already formed. Thanks to the research at [NASA](#)/GISS and

Columbia University, we may soon have a new weapon in our arsenal for predicting the areas and climate states with the greatest long-term risk for major [typhoon](#) strikes.

More information: Sovacool, B.K. 2008. The costs of failure: A preliminary assessment of major energy accidents, 1907-2007, *Energy Policy*, 36, 1802-1820, [doi:10.1016/j.enpol.2008.01.040](https://doi.org/10.1016/j.enpol.2008.01.040).

Wikipedia 2012. 2011 Pacific typhoon season. Available at en.wikipedia.org/wiki/2011_Pacific_typhoon_season. Accessed 19 Nov. 2012.

Liu, D., L. Pang, and B. Xie, 2009. Typhoon disaster in China: prediction, prevention, and mitigation. *Nat. Hazards*, 49, 421-436 [doi:10.1007/s11069-008-9262-2](https://doi.org/10.1007/s11069-008-9262-2).

Yonekura, E., and T.M. Hall, 2011. A statistical model of tropical cyclone tracks in the Western North Pacific with ENSO-dependent cyclogenesis, *J. Appl. Meteor. Climatol.*, 50, 1725-1739, [doi:10.1175/2011JAMC2617.1](https://doi.org/10.1175/2011JAMC2617.1).

Yonekura, E., and T.M. Hall, 2012. Typhoon tracks and ENSO: Separating genesis and track propagation effects, *J. Appl. Meteorol. Climatol.*, submitted.

Provided by Astrobio.net

Citation: Storms in the machine (2012, December 18) retrieved 1 May 2024 from <https://phys.org/news/2012-12-storms-machine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.