


Record diversity of ancient crustaceans may provide clues for fate of today's reef crabs

January 24 2013, by Danielle Torrent



The carapace of *Mesoparapylocheles michaeljacksoni*. The legs and tail part were not preserved. The scale bar is 1.0 mm. Credit: Neues Jahrbuch für Geologie und Paläontologie.

A University of Florida scientist has discovered a record biodiversity hotspot in Spain for 100-million-year-old crustaceans with possible


implications for present-day species living in reefs, which are declining worldwide.

Decapod crustaceans, including [crabs](#), shrimp and lobsters, are vital parts of the food chain, contributing to reef health and serving as food for many larger [marine organisms](#). Study of their early evolution may help researchers better understand how present-day species are affected by reefs declining due to changes such as [ocean acidification](#) and coral bleaching, which also threaten the animals they support.

A study published online Wednesday in [Cretaceous Research](#) reports 36 species from the abandoned Koskobilo quarry located in a fossil reef in northern Spain, including eight new species and the oldest-known spider crabs.

"The reef in Spain died soon after many decapods were still around. Something must have happened in the environment that caused reefs in the area to vanish, and with it, probably many of the decapods that were living in these reefs," said study author Adiël Klompmaker, a postdoctoral researcher at the [Florida Museum of Natural History](#) on the UF campus. "Not many decapods are known from the time after the reefs disappeared in the area."

The fauna of the locality in Spain exhibited a higher diversity than previously known, and included fossil crabs, hermit crabs and squat lobsters, said Klompmaker, who began research for the project as a graduate student at Kent State University. The fossils represent all dominant decapod groups found in reefs today, except true lobsters and shrimp, which may not have been preserved.



The carapace of one of the two oldest known spider crabs, *Cretamaja granulata*. The legs and tail part were not preserved. The scale bar is 1.0 mm. Credit: Cretaceous Research

"Today's decapods can adapt to a new environment without reefs, migrate elsewhere or go extinct," Klompmaker said. "By documenting what happened in the past, we may provide clues as to what could happen to decapod [crustaceans](#) such as crabs, shrimp and lobsters in today's deteriorating reefs."


Klompmaker researched 20 localities worldwide to determine the Koskobil quarry is the most diverse locality for decapods in the Cretaceous period 145 to 66 million years ago. The research also shows

that ancient decapods living within reefs are more diverse than those living in other parts of the ocean, which is similar to diversity patterns today.

"This shows that reefs were a popular place to feed, mate and seek shelter for decapods," Klompmaker said. "Thus, not much has changed from this perspective since the Cretaceous. Decapods still really like living in coral reefs, although today many different decapod families inhabit them."

Tropical temperatures dominated the Cretaceous period and sea levels were extremely high worldwide. The quarry, which would have been underwater at the time, is now about 35 miles from the Atlantic Ocean in northern Spain, Klompmaker said. The locality's age was determined based on ammonite fossils, ancient squid-like animals with external shells.

The study is important for understanding decapod evolution because multiple species were grouped within the same genus, rather than just one, said Francisco Vega, a paleontologist at the Universidad Nacional Autonoma de Mexico.



The carapace of the new crab *Laeviprosopon crassum*. The legs and tail part were not preserved. The scale bar is 1.0 mm. Credit: Cretaceous Research

"We would not expect such a high diversity at that time because there are very few records of crabs then," Vega said. "Also, by that time, the group was represented by so many species in a reef environment, and the number of species today is about the same, which is very significant since the diversity of the communities has prevailed ever since."

Klompaker collected fossils in the Koskobilu quarry during three field trips in 2008, 2009 and 2010 with a team of researchers from Spain, the

Netherlands and the U.S. New species were identified by analyzing the morphology, or physical characteristics of the fossils, which include three-dimensional impressions of the decapod shell.

"We went there in 2008 and in the first two hours found two new [species](#)," Klompmaker said. "That's quite amazing—it just doesn't happen every day."

More information: The scientific article can be viewed by visiting www.sciencedirect.com/science/.../S0195667112001772

Provided by University of Florida

Citation: Record diversity of ancient crustaceans may provide clues for fate of today's reef crabs (2013, January 24) retrieved 19 April 2024 from <https://phys.org/news/2013-01-diversity-ancient-crustaceans-clues-fate.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--