


# In epigenomics, location is everything: Researchers exploit gene position to test 'histone code'

January 3 2013

---



This is an X-ray micrograph of a yeast cell, *Saccharomyces cerevisiae*, as it buds before dividing. Credit: Carolyn Larabell, UC San Francisco, Lawrence Berkeley National Laboratory and the National Institute of General Medical Sciences.

In a novel use of gene knockout technology, researchers at the University of California, San Diego School of Medicine tested the same gene inserted into 90 different locations in a yeast chromosome – and discovered that while the inserted gene never altered its surrounding chromatin landscape, differences in that immediate landscape measurably affected gene activity.

The findings, published online in the Jan. 3 issue of *Cell Reports*, demonstrate that regulation of [chromatin](#) – the combination of DNA and proteins that comprise a cell's nucleus – is not governed by a uniform "[histone code](#)" but by specific interactions between chromatin and [genetic factors](#).

"One of the main challenges of epigenetics has been to get a handle on how the position of a gene in chromatin affects its expression," said senior author Trey Ideker, PhD, chief of the Division of Genetics in the School of Medicine and professor of bioengineering in UC San Diego's Jacobs School of Engineering. "And one of the major elements of that research has been to look for a histone code, a general set of rules by which histones (proteins that fold and structure DNA inside the nucleus) bind to and affect genes."

The Cell Report findings indicate that there is no singular universal code, according to Ideker. Rather, the effect of epigenetics on gene expression or activity depends not only on the particular mix of histones and other epigenetic material, but also on the identity of the gene being expressed.

To show this, the researchers exploited an overlooked feature of an existing resource. The widely-used gene knockout library for yeast, originally created to see what happens when a particular gene is missing, was built by systematically inserting the same reporter gene into different locations. Ideker and colleagues focused on this [reporter gene](#) and observed what happens to gene expression at different locations along yeast [chromosome 1](#).

"If epigenetics didn't matter – the state of histones and DNA surrounding the gene – the expression of a gene would be the same regardless of where on the chromosome that gene is positioned," said Ideker. But in every case, [gene expression](#) was measurably influenced by interaction with nearby epigenetic players.

Ideker said the work provides a new tool for more deeply exploring how and why genes function, particularly in relation to their location.

Provided by University of California - San Diego

Citation: In epigenomics, location is everything: Researchers exploit gene position to test 'histone code' (2013, January 3) retrieved 25 April 2024 from

<https://phys.org/news/2013-01-epigenomics-exploit-gene-position-histone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.