

Wood on the seafloor: An oasis for deep-sea life

January 22 2013



One of the wood experiments after one year at the seafloor. Wood-boring bivalves of the genus Xylophaga had populated the wood. Credit: Christina Bienhold, MPI for Marine Microbiology

Trees do not grow in the deep sea, nevertheless sunken pieces of wood can develop into oases for deep-sea life - at least temporarily until the wood is fully degraded. A team of Max Planck researchers from



Germany now showed how sunken wood can develop into attractive habitats for a variety of microorganisms and invertebrates. By using underwater robot technology, they confirmed their hypothesis that animals from hot and cold seeps would be attracted to the wood due to the activity of bacteria, which produce hydrogen sulfide during wood degradation.

Many of the animals thriving at <u>hydrothermal vents</u> and cold seeps require special forms of energy such as methane and <u>hydrogen sulfide</u> emerging from the ocean floor. They carry bacterial symbionts in their body, which convert the energy from these compounds into food. The vents and seeps are often separated by hundreds of kilometers of deepsea desert, with no connection between them.

For a long time it was an unsolved mystery how animals can disperse between those rare oases of energy in the deep sea. One hypothesis was that sunken <u>whale carcasses</u>, large dead algae, and also sunken woods could serve as food source and temporary habitat for deep-sea animals, but only if bacteria were able to produce methane and <u>sulfur compounds</u> from it.

To tackle this question, the team deposited wood logs on the Eastern Mediterranean seafloor at depths of 1700 meters and returned after one year to study the fauna, bacteria, and chemical microgradients.





Colonization of wood in the deep sea. Credit: Bienhold et al., PLoS ONE

"We were surprised how many animals had populated the wood already after one year. The main colonizers were wood-boring bivalves of the genus Xylophaga, also named "shipworms" after their shallow-water counterparts. The wood-boring Xylophaga essentially constitute the vanguard and prepare the habitat for other followers," Bienhold said. "But they also need assistance from bacteria, namely to make use of the cellulose from the wood, which is difficult to digest."

The team of researchers observed that the wood-boring bivalves had cut large parts of the wood into smaller chips, which were further degraded by many other organisms. This activity led to the consumption of oxygen, enabling the production of hydrogen sulfide by sulfate-reducing microorganisms. And indeed, the researchers also found a mussel, which



is typically only found at <u>cold seeps</u> or similar environments where it uses sulfur compounds as an energy source. "It is amazing to see how deep-sea bacteria can transform foreign substances such as wood to provide energy for cold-seep mussels on their journey through the deep ocean", said Antje Boetius, chief scientist of the expedition. Furthermore, the researchers discovered unknown species of deep-sea worms, which have been described by taxonomic experts in Germany and the USA. Thus, sunken woods do not only promote the dispersal of rare deep-sea animals, but also form hotspots of biodiversity at the deep seafloor.

More information: Bienhold C, Pop Ristova P, Wenzhöfer F, Dittmar T, Boetius A (2013). How Deep-Sea Wood Falls Sustain Chemosynthetic Life. *PLoS ONE* 8(1): e53590 doi:10.1371/journal.pone.0053590

Provided by Max Planck Society

Citation: Wood on the seafloor: An oasis for deep-sea life (2013, January 22) retrieved 3 May 2024 from <u>https://phys.org/news/2013-01-wood-seafloor-oasis-deep-sea-life.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.