


DNAzymes and gold nanoparticles: A colorimetric assay for diagnostics in the field

February 12 2013



Infectious diseases such as malaria and syphilis can be diagnosed rapidly and reliably in the field by using a simple test developed by Canadian scientists. The test is based on the use of DNAzymes and gold nanoparticles. As the researchers report in the journal *Angewandte Chemie*, their test allows for the sensitive detection of bacteria, viruses, and parasites.

Dangerous [infectious diseases](#) must be identified in time in order to prevent them from spreading. The DNA of pathogens is an ideal [biomarker](#) and can easily be identified by PCR. However, this is only possible if expensive laboratory equipment and trained personnel are on hand. This may not be the case in remote locations or developing nations. Alternative methods that are simple and inexpensive while also

remaining sensitive and specific are needed.

Kyryl Zagorovsky and Warren C.W. Chan at the University of Toronto (Canada) have now combined two modern technologies in a novel way: They have used DNAzymes as signal amplifiers and [gold nanoparticles](#) for detection. Gold nanoparticles (GNPs) absorb light. The wavelength of the light absorbed depends on whether the nanoparticles are separate or aggregated. The difference in color can be seen with the naked eye. A solution of individual particles appears red, whereas aggregates are blue-violet in color.

DNAzymes are synthetic [DNA molecules](#) that can enzymatically split other nucleic [acid molecules](#). The researchers separated a DNAzyme into two inactive halves that both selectively bind to a specific gene segment of the pathogen to be detected. The act of binding reunites the halves and activates them.

For their test procedure, the scientists produced two sets of GNPs that bind to two different types of [DNA strand](#), type A and type B. In addition, they synthesized a three-part "linker" made of DNA. One end of the linker is the complement to type A DNA; the second end is the complement to type B DNA. The center part is designed to be split by active DNAzymes.

In the test sample with no pathogen present, the DNAzymes remain inactive and the linkers remain intact. They bind to a GNP at each end and link the GNPs into larger aggregates, causing the solution to turn blue-violet. In contrast, if there is pathogen in the sample, the DNAzymes are activated and proceed to split the linkers. Now only the bridging parts of the linker can bind to DNA strands of the GNPs, so they cannot link the GNPs together. The solution stays red. Because every activated DNAzyme splits many linkers, it amplifies the signal.

This new type of test is simple and inexpensive; it can be made to detect every kind of pathogen, as the researchers demonstrated by detecting gonorrhea, syphilis, malaria, and hepatitis B. In a freeze-dried state, the reagents can be stored with no problem – an important requirement for use in the field.

More information: Chan, W. et al. A Plasmonic DNzyme Strategy for Point-of-Care Genetic Detection of Infectious Pathogens, *Angewandte Chemie International Edition*.
[dx.doi.org/10.1002/anie.201208715](https://doi.org/10.1002/anie.201208715)

Provided by Angewandte Chemie

Citation: DNzymes and gold nanoparticles: A colorimetric assay for diagnostics in the field (2013, February 12) retrieved 18 April 2024 from <https://phys.org/news/2013-02-dnzymes-gold-nanoparticles-colorimetric-assay.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.