

# How did early primordial cells evolve?

February 28 2013

---

Four billion years ago, soon after the planet cooled enough for life to begin, primordial cells may have replicated and divided without protein machinery or cell walls, relying instead on just a flimsy lipid membrane. New research on bacteria examines exactly how these primitive cells could have evolved without such crucial structures. While the vast majority of bacteria have cell walls, many bacteria can switch to a wall-free existence called the L-form state, which could mirror the structure of primordial cells. A study published by Cell Press February 28th in the journal *Cell* reveals how bacteria in this L-form state divide and proliferate, shedding light on how the earliest forms of cellular life may have replicated.

"The main surprise for me was how simple the mechanism was. It doesn't require any sophisticated protein-based machinery," says senior study author Jeff Errington of Newcastle University. "This makes it plausible as an explanation for how very primitive cells could have proliferated in the very early days of evolution."

The cell wall is a layered structure surrounding cells that protects them and maintains their shape. It is present in all known major bacterial lineages, and it was also probably present in the last common ancestor of bacteria. This structure is so important that it is targeted by antibiotics, and many bacteria responsible for infectious diseases can switch to the L-form state to resist antibiotics.

Perhaps the most striking change associated with the L-form state is the way that the bacteria replicate. Instead of relying on precise, complicated

cell division machinery, L-form bacteria become irregularly shaped and form cell surface bulges that pinch off to become [daughter cells](#). Although [genetic mutations](#) associated with the L-form state have been identified, little is known about the [molecular mechanisms](#) underlying L-form replication.

Addressing this question in the new study, Errington and his team identified two genetic changes required for L-form growth in bacteria. One of these mutations was necessary for the increased production of fatty acids in the cell membrane, which would be expected to increase the cell's surface area relative to its volume. Indeed, the researchers found that by artificially increasing cell surface area, they could induce L-form-like shape changes and cell division. The findings suggest that a simple biophysical change—an imbalance between surface area and volume—underlies L-form cell division.

"Our study paves the way for understanding how L-form bacteria cause disease and resist antibiotics," Errington says. "It also offers a model system for future experiments aimed at exploring the possible replication mechanisms of [primitive cells](#) that could have existed before the explosion of bacterial life on the planet nearly four billion years ago."

**More information:** *Cell*, Mercier et al.: "Excess membrane synthesis drives a primitive mode of cell proliferation."

[dx.doi.org/10.1016/j.cell.2013.01.043](https://doi.org/10.1016/j.cell.2013.01.043)

Provided by Cell Press

Citation: How did early primordial cells evolve? (2013, February 28) retrieved 24 April 2024 from <https://phys.org/news/2013-02-early-primordial-cells-evolve.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.