

Large, ancient landslides delivered preferred upstream habitats for coho salmon

February 11 2013



Joshua Roering, University of Oregon, Department of Geological Sciences, has found that coho salmon thrive and prefer in broad flat valleys with multiple small tributaries in areas formed by ancient landslides. Credit: University of Oregon

A study of the Umpqua River basin in the Oregon Coast Range helps explain natural processes behind the width of valleys and provides potentially useful details for river restoration efforts designed to improve habitats for coho salmon (*Oncorhynchus kisutch*).

Coho salmon thrive in broad, flat valleys that contain multiple auxiliary channels to the main river. These valleys formed after large [landslides](#) altered the landscape, said study co-author Joshua J. Roering, professor of [geological sciences](#) at the University of Oregon. The network of secondary channels, while often devoid of water in the dry season, fill and provide a calm, safe haven away from swiftly running currents for the fish during periods of heavy rain and storms. They now have been identified as a critical component of salmon habitat.

While [geologists](#) and [hydrologists](#) long ago documented how river-channel width and depth varies, little had been done to understand how valleys beyond the channels form, and why they may be narrow or wide, Roering said. The study—appearing online ahead of print in the journal *Geology*—combined on-the-[ground observations](#) and a remote-sensing technology known as airborne Light Detection and Ranging (LiDAR).

"There are two things that this research tells us," Roering said. "Valleys are things we can see from airplanes or from maps, and they behave, if you let them, in a very simple way. They get wider as the river gets bigger. We've all seen that. What we've learned here is that if you introduce variable geology and geomorphology—big landslides, messiness that happens in the world—the valleys tend to narrow and widen, and fish appear to love that. They seem to respond to the [heterogeneity](#) that is so inherent in most real landscapes. Nature is messy, and the fish have adapted to that."

Roering used LiDAR in previous research, including a project in which he and colleagues discovered a dry lakebed that formed some 22,500 years ago by a landslide and later drained in northern California's Eel River basin. That paper—published Nov. 22, 2011, in the *Proceedings of the National Academy of Sciences*—also connected landslides to a divergence of genetics in two present-day steelhead trout populations occupying the river.

Christine May, lead author of the new paper, said that Roering's work on landslides and her own previous observations led to the newly published research. May, a biologist at James Madison University in Harrisonburg, Va., at the time of the project was a visiting UO professor under the Meierjurgan Visiting Fellowship. While a graduate student at Oregon State University, she had seen broad valleys and productive fish habitats upstream of ancient landslides while doing fieldwork after 1996 floods.

"This observation had lingered in my mind over the years, and when Josh's study on predicting the occurrence of these large landslides on a landscape scale was published I became very excited to pursue this area of research with him," she said. "Correspondingly, my colleague Kelly Burnett with the U.S. Forest Service Pacific Northwest Research Station in Corvallis published a predictive model that mapped the location of potentially productive fish habitat in the same area as Josh mapped the large landslides."

Combing the two predictive models, May said: "It became apparent that there was a strong linkage between the occurrence of large landslides and the location of highly productive salmon habitat."

For the study, May, Roering, Burnett and L. Scott Eaton, also of James Madison University, went into the field, studying the architecture of river valleys. They used LiDAR provided by the Oregon Department of Geology and Mineral Industries, focusing on Harvey and Elk creeks—part of the Umpqua River basin where Roering previously documented ancient landslides. Researchers measured the widths of the valley using the high-resolution topographic data and showed how widths varied based on watershed-mathematical formulas used to plot and map the terrain.

To understand the activities of the salmon, researchers applied topographically derived stream-layering techniques and a formula

created by Burnett in 2007 for estimating the "intrinsic potential" of each stream to provide rearing habitats for Coho salmon.

They found that valleys downstream of landslides are narrower than would be expected, which suggested that valley constriction had subverted valley widening. Deep-seated landslides, the authors noted, caused wide valley segments to occur higher in the channel network than would otherwise be expected. Off-channel streams perched in mountain drainages, they said, increase habitat diversity and lead to a greater abundance of salmonid productivity.

"This study establishes how a valley sets its width," Roering said. "More importantly, the anomalies that make valleys look different are primarily controlled by big landslides in the Oregon Coast Range. Big events occurring thousands of years ago have had a long-running, persistent impact on the valley floor. So even if landslides sound scary or deadly, they've been active out there for long periods of time, and they make great habitat for salmon over the long term."

The findings provide fundamental insights on the geological and ecological controls on salmon habitat, May said. "I do believe the results will have broad applications for endangered species conservation and management," she added.

"Dr. Roering's explanation of how destructive natural forces can create fertile habitat answers some long-lingering questions about the geomorphology of our region," said Kimberly Andrews Espy, UO vice president for research and innovation, and dean of the graduate school. "This research helps us appreciate the richness and diversity of our state by furthering our understanding of Oregon's distinctive landscape and the plants and animals that live here."

The Umpqua River was chosen because of its "amazing uniformity,"

Roering said. "If mountains are being uplifted through tectonic forces—the rocks are going upward and erosion is wearing them down—we tend to see these two forces balancing each other. This mountain range has achieved a rough, steady state. It looks the same through time, even with a dynamic interface."

Provided by University of Oregon

Citation: Large, ancient landslides delivered preferred upstream habitats for coho salmon (2013, February 11) retrieved 19 April 2024 from <https://phys.org/news/2013-02-large-ancient-landslides-upstream-habitats.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.