

New species of mylagaulids (rodentia) found from the Miocene of northern Junggar Basin, China

February 7 2013




Fig.1 Three-dimensional pictures (3d stereo images) of right P4 of *Tschalimys ckhikvadzei* (IVPP V 8107), showing distinct dentine tract. Credit: WU Wenyu

Mylogaulids is an extinct group of Aplodontioidea (Rodentia). Most genera possess high crowned and complex cheek teeth with numerous enamel bordered lakes of different depths, which vary in size and shape with wear. Thus, it is very difficult to make taxonomic determination and assess the relationship of small samples.

Dr. WU Wenyu, Institute of [Vertebrate Paleontology](#) and [Paleoanthropology](#) (IVPP), [Chinese Academy of Sciences](#), and her collaborators applied the [CT scanning](#) technology to the study of tooth morphology of mylogaulids, and identified two genera and species, *Tschalimys ckhikvadzei* and *Simpligaulus yangi* gen. et sp. nov., from the early Middle Miocene Halamagai Formation in northern Junggar Basin of Xinjiang, China. Researchers reported in the latest issue of *Vertebrata Palasiatica* 2013 (1), providing additional information on the morphology of the mylogaulids.




Fig.2 Three dimensional pictures (3d stereo images) of left p4 of *Tschalimys ckhikvadzei* (IVPP V 17928), showing distinct dentine tract. Credit: WU Wenyu

Mylogaulid originates in the Late [Oligocene](#) and diversifies through Miocene in North America. Outside North America, mylogaulids have so far only been discovered from the early Middle Miocene of Zaisan Basin of Kazakhstan and northern Junggar Basin of China in Central Asia. *Aplodontia rufa* (mountain beaver), is the only living aplodontid and the relative of mylogaulids, inhabiting forests and densely vegetated thickets in the humid regions of western North America. They burrow in soil which is moderately firm and deep with adequate drainage.

The integrated studies indicate that the Halamagai Formation is fluvial-lacustrine sediments formed during the time coincident to the Mid-Miocene Climatic Optimum. *T. ckhikvadzei* and *S. yangi* probably inhabit the humid and warm regions with forests and densely vegetated thickets as the living mountain beaver *Aplodontia rufa* does. These mylagaulids are probably immigrants or the descendants of the immigrants from west North America.



Fig.3 Three dimensional pictures (3d stereo images) of left p4 of *Simpligaulus yangi* (IVPP V 17929), showing distinct dentine tract. Credit: WU Wenyu

"Scientists traditionally studied the mylagaulids teeth by grinding serial sections. It is essential to have sufficient material for making serial sections, thus this method is not practical for the research based on very few samples. The recently wide application of computed tomography technology in the paleontology makes it possible that our three mylagaulid teeth can be observed in detail without destruction", said first author WU Wenyu of the IVPP.

Provided by Institute of Vertebrate Paleontology and Paleoanthropology

Citation: New species of mylagaulids (rodentia) found from the Miocene of northern Junggar Basin, China (2013, February 7) retrieved 19 April 2024 from <https://phys.org/news/2013-02-species-mylagaulids-rodentia-miocene-northern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.