


Spitzer and Hubble discover strobe-like flashes in suspected binary protostar

February 7 2013



NASA's Spitzer and Hubble space telescopes have teamed up to uncover a mysterious infant star that behaves like a police strobe light. Credit: NASA, ESA, J. Muzerolle (STScI), E. Furlan (NOAO and Caltech), K. Flaherty (Univ. of Ariz./Steward Observatory), Z. Balog (Max Planck Institute for Astronomy), and R. Gutermuth (Univ. Mass. Amherst)

(Phys.org)—Two of NASA's great observatories, the Spitzer and Hubble space telescopes, have teamed up to uncover a mysterious infant star that behaves like a strobe light.


Every 25.34 days, the object, designated LRL 54361, unleashes a burst of light. Although a similar phenomenon has been observed in two other young [stellar objects](#), this is the most powerful such beacon seen to date.

The heart of the fireworks is hidden behind a dense disk and envelope of dust. Astronomers propose the light flashes are caused by periodic interactions between two newly formed stars that are binary, or gravitationally bound to each other. LRL 54361 offers insights into the early stages of star formation when lots of gas and dust is being rapidly accreted, or pulled together, to form a new [binary star](#).

Astronomers theorize the flashes are caused by material suddenly being dumped onto the growing stars, known as protostars. A blast of radiation is unleashed each time the stars get close to each other in their orbits. This phenomenon, called pulsed accretion, has been seen in later stages of [star birth](#), but never in such a young system or with such intensity and regularity.

"This [protostar](#) has such large brightness variations with a precise period that it is very difficult to explain," said James Muzerolle of the [Space Telescope Science Institute](#) in Baltimore, Md. His paper recently was published in the science journal *Nature*.

Discovered by NASA's [Spitzer Space Telescope](#), LRL 54361 is a variable object inside the star-forming region IC 348, located 950 light-years from Earth. Data from Spitzer revealed the presence of protostars. Based on statistical analysis, the two stars are estimated to be no more than a few hundred thousand years old.



This infrared image from the NASA/ESA Hubble Space Telescope shows an image of protostellar object LRL 54361 and its rich cosmic neighbourhood, a region called IC 348. The protostar, which is the bright object with fan-like beams of light coming from it, located towards the right of the image, is letting off flashes of light every 25.3 days. Credit: NASA, ESA, and J. Muzerolle (STScI)

The Spitzer [infrared data](#), collected repeatedly during a period of seven years, showed unusual outbursts in the brightness of the suspected binary protostar. Surprisingly, the outbursts recurred every 25.34 days, which is a very rare phenomenon.

Astronomers used NASA's Hubble Space Telescope to confirm the Spitzer observations and reveal the detailed stellar structure around LRL 54361. Hubble observed two cavities above and below a dusty disk. The cavities are visible by tracing light scattered off their edges.

They likely were blown out of the surrounding natal envelope of dust and gas by an outflow launched near the central stars. The disk and the envelope prevent the suspected binary star pair from being observed directly. By capturing multiple images over the course of one pulse event, the Hubble observations uncovered a spectacular movement of light away from the center of the system, an optical illusion known as a light echo.

Muzerolle and his team hypothesized the pair of stars in the center of the dust cloud move around each other in a very eccentric orbit. As the stars approach each other, dust and gas are dragged from the inner edge of a surrounding disk. The material ultimately crashes onto one or both [stars](#), which triggers a flash of light that illuminates the circumstellar dust. The system is rare because close binaries account for only a few percent of our galaxy's stellar population. This is likely a brief, transitory phase in the birth of a star system.

Muzerolle's team next plans to continue monitoring LRL 54361 using other facilities including the European Space Agency's Herschel Space Telescope. The team hopes to eventually obtain more direct measurements of the binary star and its [orbit](#).

Provided by NASA's Goddard Space Flight Center

Citation: Spitzer and Hubble discover strobe-like flashes in suspected binary protostar (2013, February 7) retrieved 21 March 2023 from <https://phys.org/news/2013-02-spitzer-hubble-strobe-like-binary-protostar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.