


Doubling estimates of light elements in the Earth's core

March 5 2013, by Karen Fox



Velocity-density plots of the samples at high pressures and temperatures. The top panel shows the velocity-density plot for hcp-Fe at both 300 K and 700 K. The dashed lines shows the linear fit, while the solid line shows the power law fit, which matches the data more closely. The bottom panel shows the velocity-density relation of both hcp-Fe and the iron-silicon alloy at 300 K.

The inner core of the Earth is the remotest area on the globe, mostly

impossible to study directly. It is an area of the planet that experiences both extremely high pressure ranging from 3,300,000 to 3,600,000 times atmospheric pressure, and extremely high temperatures somewhere from 5000 to 6000 K. One way to study this area is by recording how sound waves travel across the interior, matching these profiles to known information about how sound waves travel through candidate iron alloys, and attempting to discern which materials must be present. This method requires an understanding of how sound waves travel through the potential materials present in the core.

A team of researchers utilized APS X-rays to develop a new model of how sound waves travel through iron and iron-silicon alloys, showing for the first time that increased temperatures will affect the sound wave profile, and that sound velocity and density correlate in a non-linear way. Their results suggest that the amount of [light elements](#) in the inner core could be two times more than estimated in previous studies without considering these effects.

The researchers from the University of Texas at Austin, Argonne National Laboratory, and the Carnegie Institute of Washington studied samples of what is known as hexagonal closest-packed iron (hcp Fe), which is believed to be the high-pressure phase of iron present in the Earth's core, as well as a hcp iron-silicon alloy because silicon is one of the most likely candidate light elements in the core.

The researchers measured the compressional wave velocity (in which the wave has the same direction of vibration as its direction of travel) of the samples using high-energy inelastic x-ray scattering (HERIX) and [x-ray diffraction](#) (XRD) in a resistively-heated [diamond anvil cell](#) at XSD beamline 3-ID of the APS. They made their measurements under simultaneous high-pressure and high-temperature conditions to better simulate conditions in the Earth's core.

Previous studies had suggested that the compressional sound velocity of hcp Fe was generally linear with increased density. In contrast, by subjecting their samples to unprecedented extreme conditions, the team found that the effect of high temperature at a given density on the sound velocity of iron cannot be ignored. As the temperature increased for a given density at high pressures, the [sound waves](#) slowed down. In addition, the relationship between the sound wave velocity and increased density was not linear, but instead could be better described by an empirical power-law function with concave behavior at higher densities.

The researchers incorporated this new information into models of the Earth's core to provide new estimates of the chemical composition there. The sound wave velocities that have been observed in the core correlate to a profile of hcp-Fe with approximately 8% of silicon by weight at temperatures of 6000 K. This number represents nearly twice the amount estimated in previous studies.

The team hopes to explore additional alloys to further round out estimates of the core's composition and to explain a number of enigmatic behaviors of seismic waves in this most extreme region of the planet. Also, since this study observed temperatures and pressures that are still much lower than those of the inner core, these scientists hope future studies will push the experimental conditions even further. With direct measurements of compressional wave velocities at relevant pressure and temperature conditions on the horizon, such studies may eventually answer the long-standing question of the composition of the Earth's core.

More information: Mao, Z. et al. Sound velocities of Fe and Fe-Si alloy in the Earth's core, *Proc. Natl. Acad. Sci.* 109(26), 10239 (June 26, 2012). [DOI:10.1073/pnas.1207086109](https://doi.org/10.1073/pnas.1207086109)

Provided by Argonne National Laboratory

Citation: Doubling estimates of light elements in the Earth's core (2013, March 5) retrieved 3 May 2024 from <https://phys.org/news/2013-03-elements-earth-core.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.