

Sequencing without PCR reduces bias in measuring biodiversity

March 26 2013

DNA barcode sequencing without the amplification of DNA by PCR beats the problem of false positives which can inflate estimates of biodiversity, finds a study published in BioMed Central and BGI Shenzhen's open access journal *GigaScience*. This method tested on a bulk 'squashome' of mixed insect samples is also able rapidly and cost-effectively estimate biomass.

Often samples collected in the field are too small be sequenced directly. Traditionally, to get round this problem, DNA from the sample is amplified using multiple cycles of PCR. However this process is highly dependent on the chemical and physical properties of each piece of DNA and consequently this approach can selectively amplify certain sequences more than others.

Researchers from BGI and the China National [GeneBank](#)-Shenzhen have developed a method of COI metabarcoding which uses mitochondrial enrichment prior to [DNA extraction](#), rather than PCR, to ensure a high enough concentration of DNA for next generation ultra-deep sequencing to be accurate.

Successfully tested on 69 [arthropods](#) collected from a mountain in sub-tropical China this method was able to correctly match insects to their reference samples without introducing the false positives often seen using traditional methods. False positives are the main culprit behind biodiversity inflation. A few samples from insects with a body length of less than 5mm contained too little material for this method to work.

The screen also managed to identify bacteria within the samples, two of which were *Wolbachia*, a [symbiont](#) frequently found in insects, plus Legionellaceae and Bartonellaceae which can cause disease in humans. Additionally DNA was found from a member of the Lepidoptera order (butterflies and moths) even though it was not in the original insect mixture. It is thought that this probably either came from the gut of an insect that ate it, or from an undetected fragment or egg.

Commenting on the ability of this study in evaluating biodiversity Dr Xin Zhou who led this study commented, "We also found that sequencing 'volume' (nucleotide numbers) strongly indicated total biomass for the species in a bulk sample. By being able to identify both species and the biomass for each species more accurately than traditional PCR-based sequencing, this method could revolutionise biodiversity research and biomonitoring."

More information: Ultra-deep sequencing enables high-fidelity recovery of biodiversity for bulk arthropod samples without PCR amplification, Xin Zhou, Yiyuan Li, Shanlin Liu, Qing Yang, Xu Su, Lili Zhou, Min Tang, Ribei Fu, Jiguang Li and Quanfei Huang *GigaScience* 2013, 2:4 [doi:10.1186/2047-217X-2-4](https://doi.org/10.1186/2047-217X-2-4)

NGS biodiversity data

Zhou, X; Li, Y; Liu, S; Yang, Q; Su, X; Zhou, L; Tang, M; Fu, R; Li, J
GigaScience Database 2013 dx.doi.org/10.5524/100045

NGS Biodiversity software

Zhou, X; Li, Y; Liu, S; Yang, Q; Su, X; Zhou, L; Tang, M; Fu, R; Li, J;
Huang, Q, *GigaScience* Database 2013 dx.doi.org/10.5524/100046
(Provides software and supporting material)

Provided by BioMed Central

Citation: Sequencing without PCR reduces bias in measuring biodiversity (2013, March 26)
retrieved 26 June 2024 from <https://phys.org/news/2013-03-sequencing-pcr-bias-biodiversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.