

Summer melt season is getting longer on the Antarctic Peninsula, research shows

March 27 2013

New research from the Antarctic Peninsula shows that the summer melt season has been getting longer over the last 60 years. Increased summer melting has been linked to the rapid break-up of ice shelves in the area and rising sea level.

The [Antarctic Peninsula](#) – a mountainous region extending northwards towards South America – is warming much faster than the rest of Antarctica. Temperatures have risen by up to 3 oC since the 1950s – three times more than the global average. This is a result of a strengthening of local [westerly winds](#), causing warmer air from the sea to be pushed up and over the peninsula. In contrast to much of the rest of Antarctica, [summer temperatures](#) are high enough for snow to melt.

This summer melting may have important effects. Meltwater may enlarge cracks in [floating ice](#) shelves which can contribute to their retreat or collapse. As a result, the speed at which glaciers flow towards the sea will be increased. Also, melting and refreezing causes snow layers to become thinner and more dense, affecting the height of the snow surface above sea level. Scientists need to know this so they can interpret satellite data correctly.

Dr Nick Barrand, who carried out the research while working for the [British Antarctic Survey](#), led an analysis of data from 30 weather stations on the peninsula. "We found a significant increase in the length of the melting season at most of the stations with the longest temperature records" he says. "At one station the average length of the melt season

almost doubled between 1948 and 2011."

To build up a more complete picture across the whole peninsula, the team (funded by the European Union's ice2sea programme) also analysed satellite data collected by an instrument called a scatterometer. Using microwave reflections from the ice sheet surface, the [scatterometer](#) was able to detect the presence of meltwater. The team were able to produce maps of how the melt season varied from 1999 to 2009, and showed that several major ice shelf breakup events coincided with longer than usual melt seasons. This supports the theory that enlargement of cracks by [meltwater](#) is the main mechanism for ice shelf weakening and collapse.

The researchers also compared data from both the satellite and [weather stations](#) with the output of a state-of-the-art regional climate model.

Dr Barrand, who now works at the University of Birmingham, says, "We found that the model was very good at reproducing the pattern and timing of the melt, and changes in melting between years. This increases confidence in the use of climate models to predict future changes to snow and ice cover in the Antarctic Peninsula."

More information: Trends in Antarctic Peninsula surface melting conditions from observations and regional climate modeling will be officially published in the *Journal of Geophysical Research* this week.

Provided by British Antarctic Survey

Citation: Summer melt season is getting longer on the Antarctic Peninsula, research shows (2013, March 27) retrieved 13 May 2024 from <https://phys.org/news/2013-03-summer-season-longer-antarctic-peninsula.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.