

# Astronomer studies far-off worlds through 'characterization by proxy'

April 26 2013, by Peter Kelley

---

(Phys.org) —A University of Washington astronomer is using Earth's interstellar neighbors to learn the nature of certain stars too far away to be directly measured or observed, and the planets they may host.

"Characterization by proxy" is the technique used by Sarah Ballard, a post-doctoral researcher at the UW, to infer the properties of small, relatively cool stars too distant for measurement, by comparing them to closer stars that now can be directly observed.

Ballard is lead author of a study accepted for publication in *The Astrophysical Journal* that used this method and observations from the Kepler [Space Telescope](#) to learn the nature of the [distant star](#) Kepler-61.

Our understanding of the size and temperature of planets depends crucially on the size and temperature of the stars they orbit. Astronomers already have a robust way to discern the physical properties of solar-type stars—those like the sun—by measuring the light they emit at different wavelengths and matching that to synthetically created spectra.

"The challenge is that small stars are incredibly difficult to characterize," Ballard said. Those theoretical methods don't work well for what are called M-dwarf stars, lower-mass stars about half the size of the sun and smaller—which is too bad, because such stars make up about three-quarters of the universe.

Ballard is using the characterization by proxy method to try to fill this

[knowledge gap](#). She is building on what she calls "truly remarkable" work by astronomer Tabettha Boyajian, now at Yale University, who uses a near-infrared interferometer—an array of telescopes working in unison studying light wavelengths a bit longer than visible light—to resolve the physical size of relatively [nearby stars](#).

Ballard said her characterization by proxy method takes "full advantage that there now exists this precious sample" of relatively nearby stars that have been directly measured. You could say the method borrows a bit from Greek mathematician Euclid, whose first "common notion" held that things that equal the same thing must necessarily also equal each other.

In the new paper, Ballard and co-authors used this reasoning to learn about Kepler-61b, a planet orbiting near the inner edge of the [habitable zone](#) of the distant, low-mass star Kepler-61, about 900 light-years away in the Cygnus Constellation. A star's habitable zone is that swath of space just right for an orbiting planet's surface water to be in liquid form, thus giving life a chance.

She did this by comparing it to temperature size averages from four spectroscopically similar stars between 12 and 25 light-years away in the Ursa Major and Cygnus constellations. A light-year is about 6 trillion miles.

A funny thing also happened along the way: Kepler-61 turned out to be bigger and hotter than expected, which in turn recalibrated planet [Kepler -61b](#)'s relative size upward as well—meaning it, too, would be hotter than previously thought and no longer a resident of the star's habitable zone.

All of this caused Ballard to informally subtitle the paper, "How Nearby [Stars](#) Bumped a Planet out of the Habitable Zone."

**More information:** [iopscience.iop.org/0004-637X](https://iopscience.iop.org/0004-637X)

Provided by University of Washington

Citation: Astronomer studies far-off worlds through 'characterization by proxy' (2013, April 26)  
retrieved 23 April 2024 from

<https://phys.org/news/2013-04-astronomer-far-off-worlds-characterization-proxy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.