

Bird fossil sheds light on how swift and hummingbird flight came to be

May 1 2013



12 centimeters from head to tail, *E. rowei* was an evolutionary precursor to the group that includes today's swifts and hummingbirds. Credit: Photo contributed by Lance Grande of the Field Museum of Natural History



A tiny bird fossil discovered in Wyoming offers clues to the precursors of swift and hummingbird wings. The fossil is unusual in having exceptionally well-preserved feathers, which allowed the researchers to reconstruct the size and shape of the bird's wings in ways not possible with bones alone.

Researchers spotted the specimen—the nearly complete skeleton of a bird that would have fit in the palm of your hand and weighed less than an ounce—while working at the Field Museum of Natural History in Chicago.

The newly discovered bird was named *Eocypselus rowei*, in honor of John W. Rowe, Chairman of the Field Museum's Board of Trustees.

First collected in southwestern Wyoming in a fossil site known as the Green River Formation, *E. rowei* lived roughly 50 million years ago, after the dinosaurs disappeared but before the earliest humans came to be.

E. rowei was a tiny bird – only twelve centimeters from head to tail. Feathers account for more than half of the bird's total wing length.

To find out where the fossil fit in the bird family tree, the researchers compared the specimen to extinct and modern day species. Their analyses suggest that the bird was an evolutionary precursor to the group that includes today's swifts and hummingbirds.

Given the differences in <u>wing shape</u> between these two closely related groups of <u>birds</u>, scientists have puzzled over how swift and hummingbird flight came to be. Finding fossil relatives like this specimen is key to figuring that out, the researchers say.

"This <u>fossil</u> bird represents the closest we've gotten to the point where



swifts and hummingbirds went their separate ways," said lead author Daniel Ksepka of the National <u>Evolutionary Synthesis</u> Center in Durham, North Carolina.

Hummingbirds have short wings relative to their bodies, which makes them good at hovering in mid-air. Swifts have super-long wings for gliding and high-speed flight. But the wings of *E. rowei* were somewhere in between.

"[Based on its wing shape] it probably wasn't a hoverer, like a hummingbird, and it probably wasn't as efficient at fast flight as a swift," Ksepka said.

The shape of the bird's wings, coupled with its tiny size, suggest that the ancestors of today's swifts and hummingbirds got small before each group's unique flight behavior came to be. "<u>Hummingbirds</u> came from small-bodied ancestors, but the ability to hover didn't come to be until later," Ksepka explained.

Closer study of the feathers under a scanning electron microscope revealed that carbon residues in the fossils—once thought to be traces of bacteria that fed on feathers—are fossilized melanosomes, tiny cell structures containing melanin pigments that give birds and other animals their color. The findings suggest that the ancient bird was probably black and may have had a glossy or iridescent sheen, like swifts living today. Based on its beak shape it probably ate insects, the researchers say.

The other authors of this study were Julia Clarke, Sterling Nesbitt and Felicia Kulp of the University of Texas at Austin, and Lance Grande of the Field Museum of Natural History.

The results will appear in the May 1 issue of the journal *Proceedings of the Royal Society B*.



More information: Ksepka, D., et al. (2013). "Fossil evidence of wing shape in a stem relative of swifts and hummingbirds (Aves, Pan-Apodiformes)." *Proceedings of the Royal Society B*.

Provided by National Evolutionary Synthesis Center (NESCent)

Citation: Bird fossil sheds light on how swift and hummingbird flight came to be (2013, May 1) retrieved 2 May 2024 from <u>https://phys.org/news/2013-04-bird-fossil-swift-hummingbird-flight.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.