


New coating could enable major boost in solar-cell efficiency

April 18 2013, by David Chandler



Credit: Christine Daniloff/MIT

Throughout decades of research on solar cells, one formula has been considered an absolute limit to the efficiency of such devices in converting sunlight into electricity: Called the Shockley-Queisser efficiency limit, it posits that the ultimate conversion efficiency can never exceed 34 percent for a single optimized semiconductor junction.



Now, researchers at MIT have shown that there is a way to blow past that limit as easily as today's jet fighters zoom through the <u>sound barrier</u> —which was also once seen as an ultimate limit.

Their work appears this week in a report in the journal *Science*, coauthored by graduate students including Daniel Congreve, Nicholas Thompson, Eric Hontz and Shane Yost, alumna Jiye Lee '12, and professors Marc Baldo and Troy Van Voorhis.

The principle behind the barrier-busting technique has been known theoretically since the 1960s, says Baldo, a professor of electrical engineering at MIT. But it was a somewhat obscure idea that nobody had succeeded in putting into practice. The MIT team was able, for the first time, to perform a successful "proof of principle" of the idea, which is known as singlet exciton <u>fission</u>. (An exciton is the <u>excited state</u> of a molecule after absorbing energy from a photon.)

In a standard <u>photovoltaic</u> (PV) cell, each photon knocks loose exactly one electron inside the PV material. That loose electron then can be harnessed through wires to provide an electrical current.

But in the new technique, each photon can instead knock two electrons loose. This makes the process much more efficient: In a standard cell, any <u>excess energy</u> carried by a photon is wasted as heat, whereas in the new system the extra energy goes into producing two electrons instead of one.

While others have previously "split" a photon's energy, they have done so using ultraviolet light, a relatively minor component of sunlight at Earth's surface. The new work represents the first time this feat has been accomplished with visible light, laying a pathway for practical applications in solar PV panels.



This was accomplished using an organic compound called pentacene in an organic solar cell. While that material's ability to produce two excitons from one photon had been known, nobody had previously been able to incorporate it within a PV device that generated more than one electron per photon.

"Our whole project was directed at showing that this splitting process was effective," says Baldo, who is also the director of the Center for Excitonics, sponsored by the U.S. Department of Energy. "We showed that we could get through that barrier."

The theoretical basis for this work was laid long ago, says Congreve, but nobody had been able to realize it in a real, functioning system. "In this system," he says, "everyone knew you could, they were just waiting for someone to do it."

Since this was just a first proof of principle, the team has not yet optimized the energy-<u>conversion efficiency</u> of the system, which remains less than 2 percent. But ratcheting up that efficiency through further optimization should be a straightforward process, the researchers say. "There appears to be no fundamental barrier," Thompson says.

While today's commercial solar panels typically have an efficiency of at most 25 percent, a silicon solar cell harnessing singlet fission should make it feasible to achieve <u>efficiency</u> of more than 30 percent, Baldo says—a huge leap in a field typically marked by slow, incremental progress. In solar cell research, he notes, people are striving "for an increase of a tenth of a percent."

Solar panel efficiencies can also be improved by stacking different solar cells together, but combining solar cells is expensive with conventional solar-cell materials. The new technology instead promises to work as an inexpensive coating on <u>solar cells</u>.



The work made use of a known material, but the team is now exploring new materials that might perform the same trick even better. "The field is working on materials that were chanced upon," Baldo says—but now that the principles are better understood, researchers can begin exploring possible alternatives in a more systematic way.

More information: "External Quantum Efficiency Above 100% in a Singlet-Exciton-Fission–Based Organic Photovoltaic Cell," by D.N. Congreve, *Science*, 2013.

Provided by Massachusetts Institute of Technology

Citation: New coating could enable major boost in solar-cell efficiency (2013, April 18) retrieved 25 April 2024 from <u>https://phys.org/news/2013-04-coating-enable-major-boost-solar-cell.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.