

# Research sheds light on how patterns form in bird feathers

April 26 2013, by Lin Edwards

---



Feathers exhibit complex pigment patterns and can be a great model to learn how morphogenesis patterns stem cells into organized tissues. Lin et al. identify feather follicle melanocyte progenitors and demonstrate how complex patterns can emerge from multi-dimensional modulations of simple regulatory mechanisms. Feathers shown are from Silver Laced Wyandotte, Barred Rock, and Guineafowl (from left to right). Credit: Ting Xin Jiang. Figure design: Randall B Wideltiz

(Phys.org) —A new study by scientists in the U.S. and Taiwan has

shown that birds have colorful and patterned feathers because of specific cellular interactions involving stem cells rather than through the direct involvement of encoding in DNA.

Until now the patterning of pigmentation in bird feathers has been poorly understood, but the patterns and colors are known to be important for attracting mates, protecting from predators, and they also affect [bird behavior](#) and can exhibit [seasonal variations](#). The new study found that the complex patterns formed by the pigments in the feather follicles were the result of several distinct mechanisms, including changes in melanocyte formation.

The research team found that the first important process was the production of [melanocytes](#), which are also found in hair, where they produce the [pigment melanin](#). In hair, melanocyte stem cells (McSCs) are found in the follicle bulge at the base of the follicle. The new research found melanocyte progenitor [stem cells](#) in a ring in the proximal follicle, and showed that the melanocytes distribute upwards into the epithelial cylinder of the feather as it grows. The different patterns in each feather are determined by the arrangement and differentiation of pigmented and unpigmented melanocytes. They found the melanocyte distribution could differ from the front to the back of the feather, and from the center to the outer edge. The patterns and colors could also change over time as the feather grew.

The research team, led by Dr. C.M. Chuong of the Keck School of Medicine at the University of Southern California and the National Taiwan University, looked at the processes involved in pigment formation in several [species of birds](#), including chickens. They first mapped the [progenitor cells](#) in regenerating feather follicles, and found that during the growth phase the progenitors gave rise to both pigmented and unpigmented melanocytes. During the resting phase the progenitor cells were quiescent.



Exhibition chickens displaying distinct black and white pigment patterns created by spatial and temporal control of melanocyte distribution and differentiation. From top left, male Silver Spangled Hamburg; male Barred Wyandotte; female Silver Laced Wyandotte; male Silver laced Wyandotte. Credits: Male Silver Spangled Hamburg: Jim Legendre; Male Barred Wyandotte: Grant Brereton; Female Silver Laced Wyandotte and male Silver laced Wyandotte: Doug and Pete Akers

They next transplanted melanocyte progenitor cells from adult quail feathers into chicken embryos and found that the chickens produced pigmented feathers.

The team looked at white feathers from several types of white chicken to determine whether the whiteness was caused by the absence of

melanocytes or suppressed melanocyte differentiation, and they examined feathers with different colors at each end, and striped feathers, to try to determine how these patterns were formed. They found that while melanocytes may be present in unpigmented regions and there were still melanocyte progenitor cells, the melanocytes were not differentiated.

The researchers also found that the patterning was affected by the topological shape of the melanocyte stem cell progenitor niche. The appearance of pigment was also affected by agouti, a peptide found in the pulp in the feather follicles. Agouti had a modifying effect on pigmentation, since it was found to inhibit melanocytes and produce unpigmented regions.

**More information:** Topology of Feather Melanocyte Progenitor Niche Allows Complex Pigment Patterns to Emerge, *Science* [DOI: 10.1126/science.1230374](https://doi.org/10.1126/science.1230374)

## **ABSTRACT**

Color patterns of bird plumage affect animal behavior and speciation. Diverse patterns are present in different species and within the individual. Here, we study the cellular and molecular basis of feather pigment pattern formation. Melanocyte progenitors are distributed as a horizontal ring in the proximal follicle, sending melanocytes vertically up into the epithelial cylinder which gradually emerges as feathers grow. Different pigment patterns form by modulating the presence, arrangement, or differentiation of melanocytes. A layer of peripheral pulp further regulates pigmentation via patterned agouti expression. Lifetime feather cyclic regeneration resets pigment patterns for physiological needs. Thus, the evolution of stem cell niche topology allows complex pigment patterning via combinatorial co-option of simple regulatory mechanisms.

© 2013 Phys.org

Citation: Research sheds light on how patterns form in bird feathers (2013, April 26) retrieved 19 April 2024 from <https://phys.org/news/2013-04-patterns-bird-feathers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.