

Scientists reach the ultimate goal: Controlling chirality in carbon nanotubes

April 29 2013



An ultimate goal in the field of carbon nanotube research is to synthesise single-walled carbon nanotubes (SWNTs) with controlled chiralities. Twenty years after the discovery of SWNTs, scientists from Aalto University in Finland, A.M. Prokhorov General Physics Institute RAS in Russia and the Center for Electron Nanoscopy of Technical University of Denmark (DTU) have managed to control chirality in carbon nanotubes during their chemical vapor deposition synthesis

<u>Carbon nanotube</u> structure is defined by a pair of integers known as chiral indices (n,m), in other words, chirality.



"<u>Chirality</u> defines the optical and <u>electronic properties</u> of carbon nanotubes, so controlling it is a key to exploiting their practical applications," says Professor Esko I. Kauppinen, the leader of the Nanomaterials Group in Aalto University School of Science.

Over the years, substantial progress has been made to develop various structure-controlled synthesis methods. However, precise control over the chiral structure of SWNTs has been largely hindered by a lack of practical means to direct the formation of the metal nanoparticle catalysts and their catalytic dynamics during tube growth.

"We achieved an epitaxial formation of Co nanoparticles by reducing a well-developed solid solution in CO," reveals Maoshuai He, a <u>postdoctoral researcher</u> at Aalto University School of <u>Chemical</u> <u>Technology</u>.

"For the first time, the new catalyst was employed for selective growth of SWNTs," adds senior staff scientist Hua Jiang from Aalto University School of Science.

By introducing the new catalysts into a conventional CVD reactor, the research team demonstrated preferential growth of semiconducting SWNTs (290%) with an exceptionally high population of (6,5) tubes (53%) at 500 °C. Furthermore, they also showed a shift of the chiral preference from (6,5) tubes at 500 °C to (7, 6) and (9, 4) nanotubes at 400 °C.

"These findings open new perspectives both for structural control of SWNTs and for elucidating their growth mechanisms, thus are important for the fundamental understanding of science behind nanotube growth," comments Professor Juha Lehtonen from Aalto University.

The research has been recently published in a new Nature Publishing



Group journal Scientific Reports, 3 (2013), 1460.

More information: Link to article: <u>www.nature.com/srep/2013/13031</u> ... /full/srep01460.html

Provided by Aalto University

Citation: Scientists reach the ultimate goal: Controlling chirality in carbon nanotubes (2013, April 29) retrieved 1 May 2024 from <u>https://phys.org/news/2013-04-scientists-ultimate-goal-chirality-carbon.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.