

# Wild mice have natural protection against Lyme borreliosis

April 4 2013



Ticks suck blood. About the saliva of infected ticks, *Borrelia* get into the blood stream of mice. Credit: UZH

Like humans, mice can become infected with *Borrelia*. However, not all mice that come into contact with these bacteria contract the dreaded Lyme disease: Animals with a particular gene variant are immune to the

bacteria, as scientists from the universities of Zurich and Lund demonstrate. Wild mice are the primary hosts for *Borrelia*, which are transmitted by ticks.

Springtime spells tick-time. Lyme [borreliosis](#) is the most common tick-borne disease in Switzerland: around 10,000 people a year become infected with the pathogen. The actual hosts for *Borrelia*, however, are wild mice. Like in humans, the pathogen is also transmitted by ticks in mice. Interestingly, not all mice are equally susceptible to the bacterium and individual animals are immune to the pathogen. Scientists from the universities of Zurich and Lund headed by [evolutionary biologist](#) Barbara Tschirren reveal that the difference in vulnerability among the animals is genetic in origin.

## **Protective gene variant**

Tschirren and colleagues examined wild mice for signs of a *Borrelia* infection in a large-scale field study. *Borrelia afzelii* – the scientific name for the bacteria – feed on mouse blood. The researchers discovered that mice with a particular variant of the antigen receptor TLR2 were around three times less susceptible to *Borrelia*. "The immune system of mice with this receptor variant recognizes the pathogen better and can trigger an immune response more quickly to destroy the *Borrelia* in time," says Tschirren. Infected mice exhibit similar symptoms to humans – especially joint complaints. Consequently, in the wild infected mice probably do not survive for very long and weakened animals soon fall victim to foxes and birds of prey.

## **Arms race between mice and *Borrelia***

The protective [gene variant](#) is advantageous for its carriers and, according to the researchers, gradually becoming prevalent in the [mouse](#)

[population](#). Nonetheless, it is unlikely that all mice will one day be resistant to *Borrelia*. "The increasing resistance in the host is bound to lead to adaptations in *Borrelia*," predicts Tschirren. "We can observe the evolutionary adaptation through the rearmament in mice and the pathogen."

People also have the [antigen receptor](#) TLR2, but not the resistant gene variant observed in mice. Whether the evolutionary arms race between mice and *Borrelia* will have repercussions for people remains to be seen. According to Tschirren, the bacterium does not necessarily have to become more aggressive for humans.

Tschirren, B. et al. Polymorphisms at the innate immune receptor TLR2 are associated with *Borrelia* infection in a wild rodent population. *Proceedings of the Royal Society B*, 20130364. April 3 , 2013. doi: 10.1098/rspb.2013.0364

Provided by University of Zurich

Citation: Wild mice have natural protection against Lyme borreliosis (2013, April 4) retrieved 9 April 2024 from <https://phys.org/news/2013-04-wild-mice-natural-lyme-borreliosis.html>

|  |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|