


Competition in the quantum world

May 20 2013



The physicists engineer a classical environment, which generates dissipative dynamics, leading to fragile long-range quantum mechanical correlations between distant particles. Credit: IQOQI/Ritsch

Innsbruck physicists led by Rainer Blatt and Peter Zoller experimentally gained a deep insight into the nature of quantum mechanical phase transitions. They are the first scientists that simulated the competition between two rival dynamical processes at a novel type of transition

between two quantum mechanical orders. They have published the results of their work in the journal *Nature Physics*.

"When water boils, its molecules are released as vapor. We call this change of the physical state of matter a phase transition," explains Sebastian Diehl from the Institute of Theoretical Physics at the University of Innsbruck. Together with his colleagues from the Institute for [Experimental Physics](#) and the theorist Markus Mueller from the Complutense University of Madrid, he studied the transition between two quantum mechanical orders in a way never before observed. The [quantum physicists](#) in Innsbruck use a new device for the experiment, which is currently considered to be one of the most promising developments in [quantum physics](#): a quantum simulator. It is based on a small-scale quantum computer and can simulate [physical phenomena](#) a [classical computer](#) cannot investigate efficiently. "Such a quantum simulator allows us to experimentally study [quantum phenomena](#) in many-body systems that are coupled to their environment," explain experimental physicists Philipp Schindler and Thomas Monz.

Observing the competition

With just a few trapped ions the scientists simulate the complex physical processes of quantum mechanical phase transitions. To achieve this, they have to manipulate and control the particles with high accuracy; the experimental physicists in Innsbruck are world leaders in this field. "For this experiment we use a programmable quantum simulator with up to five ions," says Philipp Schindler. One of the particles is used as a means to couple the system to the classical environment in a controlled manner. The other ions are used for carrying out quantum operations. "We call this an open [quantum simulator](#). Usually we want to suppress this coupling because it destroys the fragile quantum effects in the system. Here, however, we use it to bring order into the quantum mechanical system," explains Schindler. "In our specific case, we engineer a classical

environment, which generates dissipative dynamics, leading to fragile long-range quantum mechanical correlations between distant particles." In the following step, this dynamics is then set in competition with a different type of interactions, which interrupts the dynamics that create the quantum mechanical order. "By doing this, we are able to observe how the competition between these two processes takes place and what precisely occurs right at the transition between two distinct orders of matter," explains theoretical physicist Sebastian Diehl.

Error reduction

The experiment demands an enormous degree of precision, which requires immediate error corrections to be able to simulate the physical processes correctly. Since a comprehensive error correction, as developed for quantum computers, involves considerable resource overheads, the physicists in Innsbruck chose another promising alternative path. They identified the most important sources of error occurring during the simulation and specifically targeted them. Schindler is convinced: "This way of error reduction will surely set an example for other experiments. While general quantum error correction remains a long-term goal, we may be able to successfully use this type of error correction a lot sooner for reliable quantum simulation of larger systems," adds Markus Mueller.

Interweaving theory with experiment

Such an experimental study of the nature of quantum mechanical [phase transitions](#) is internationally unique. It was only possible because advanced experimental know-how was successfully combined with theoretical research, which was carried out in close collaboration between physicists from Innsbruck and Madrid. "This link between theoretical and [experimental physicists](#) who work closely together, and

in Innsbruck under one roof, is possible in very few places. It is also one of the great strengths of quantum physics research carried out in Innsbruck. And this research, once again, led us into an area of physics that hadn't been explored before," says Rainer Blatt. "In this experiment the physics of many-body systems is successfully simulated with a few trapped ions. This clearly shows the potential and the possibilities of quantum simulation," adds Peter Zoller.

More information: Quantum simulation of dynamical maps with trapped ions. P. Schindler, M. Müller, D. Nigg, J. T. Barreiro, E. A. Martinez, M. Hennrich, T. Monz, S. Diehl, P. Zoller und R. Blatt, Advance online publication, *Nature Physics* on May 19, 2013, [DOI: 10.1038/NPHYS2630](https://doi.org/10.1038/NPHYS2630)

Provided by University of Innsbruck

Citation: Competition in the quantum world (2013, May 20) retrieved 3 May 2024 from <https://phys.org/news/2013-05-competition-quantum-world.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--