


Gold nanocrystal vibration captured on billion-frames-per-second film (w/ video)

May 23 2013



The acoustic phonons can be visualized on the surface as regions of contraction (blue) and expansion (red). Also shown are two-dimensional images comparing the experimental results with theory and molecular dynamics simulation. The scale bar is 100 nanometers. Credit: Jesse Clark/UCL

A billion-frames-per-second film has captured the vibrations of gold nanocrystals in stunning detail for the first time.

The film, which was made using 3D imaging pioneered at the London Centre for Nanotechnology (LCN) at UCL, reveals important

information about the composition of gold. The findings are published in the journal *Science*.

Jesse Clark, from the LCN and lead author of the paper said: "Just as the sound quality of a musical instrument can provide great detail about its construction, so too can the vibrations seen in materials provide important information about their composition and functions."

"It is absolutely amazing that we are able to capture snapshots of these nanoscale motions and create movies of these processes. This information is crucial to understanding the response of materials after [perturbation](#)."

Scientists found that the vibrations were unusual because they start off at exactly the same moment everywhere inside the crystal. It was previously expected that the effects of the excitation would travel across the gold nanocrystal at the speed of sound, but they were found to be much faster, i.e., supersonic.

The new images support [theoretical models](#) for [light interaction](#) with metals, where energy is first transferred to electrons, which are able to short-circuit the much slower motion of the atoms.

The team carried out the experiments at the SLAC National Accelerator Laboratory using a revolutionary X-ray laser called the "[Linac Coherent Light Source](#)". The pulses of X-rays are extremely short (measured in [femtoseconds](#), or quadrillionths of a second), meaning they are able to freeze all motion of the atoms in any sample, leaving only the electrons still moving.

However, the X-ray pulses are intense enough that the team was able to take single snapshots of the vibrations of the gold [nanocrystals](#) they were examining. The [vibration](#) was started with a short pulse of [infrared light](#).

The vibrations were imaged a short time later in 3D using the coherent diffraction imaging methods pioneered in LCN by the Robinson group. The 3D movies reveal in exquisite detail the distortions taking place within the nanocrystal, with the fastest vibrations repeating every 90 picoseconds.

Professor Robinson, also from the LCN and the group leader, said: "This work represents an impressive example of teamwork by about a hundred people at SLAC. The SLAC linear accelerator was built in 1957 in direct response to the news of Sputnik.

"After compelling 50 years of sensational high energy physics, that machine has been refitted as a laser by the addition of a 100m long array of magnets. This 3km-sized machine produces a beam which is focused onto a crystal smaller than a micron in a pulse so short that all motion of its atoms is frozen still."

More information: 'Ultrafast three dimensional imaging of lattice dynamics in gold nanocrystals' is published online today in the journal *Science*.

Provided by University College London

Citation: Gold nanocrystal vibration captured on billion-frames-per-second film (w/ video) (2013, May 23) retrieved 4 May 2024 from <https://phys.org/news/2013-05-gold-nanocrystal-vibration-captured-billion-frames-per-second.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.