

# National study of nanomaterial toxicity sets stage for policies to address health risks

May 6 2013

---

For the first time, researchers from institutions around the country have conducted an identical series of toxicology tests evaluating lung-related health impacts associated with widely used engineered nanomaterials (ENMs). The study provides comparable health risk data from multiple labs, which should help regulators develop policies to protect workers and consumers who come into contact with ENMs.

Researchers have done a great deal of toxicological research on ENMs over the past 10 years, but the results have often been difficult to interpret. This is because ENMs from different sources had different chemical and physical properties, and because investigators used different protocols to conduct the experiments.

"The goal of creating this multicenter consortium was to have multiple labs recreate key studies using the same materials and protocols, so that policy-makers have access to consistent, comparable results from multiple institutions," says Dr. James Bonner, an associate professor of environmental and [molecular toxicology](#) at NC State and lead author of a paper describing the work.

For this study, researchers from eight institutions used mouse and rat models to look at pulmonary health effects related to exposure to titanium dioxide nanoparticles and carbon nanotubes.

The researchers found that carbon nanotubes, which are used in everything from bicycle frames to high performance electronics,

produced inflammation and inflammatory lesions in the lower portions of the lung. However, the researchers found that the nanotubes could be made less hazardous if treated to remove excess [metal catalysts](#) used in the manufacturing process or modified by adding carboxyl groups to the [outer shell](#) of the tubes to make them more easily dispersed in [biological fluids](#).

The researchers also found that [titanium dioxide](#) nanoparticles also caused inflammation in the lower regions of the lung. Belt-shaped titanium nanoparticles caused more cellular damage in the lungs, and more pronounced lesions, than spherical nanoparticles.

"The findings are significant, but the real take-away message here is that the multicenter consortium concept works – and that means this is a starting point for assessing nanomaterials using this approach," Bonner says. "I'm optimistic that this will serve as a blueprint for similar efforts, which will give regulators comparable data across institutions that will be easier for them to interpret."

**More information:** The paper, "Inter-laboratory Evaluation of Rodent Pulmonary Responses to Engineered Nanomaterials," was published May 6 in *Environmental Health Perspectives*.

Provided by North Carolina State University

Citation: National study of nanomaterial toxicity sets stage for policies to address health risks (2013, May 6) retrieved 18 April 2024 from <https://phys.org/news/2013-05-national-nanomaterial-toxicity-stage-policies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.