


## Researchers develop unique method for creating uniform nanoparticles

May 6 2013, by Rick Kubetz

---




This is a micrograph showing array of atoms and tetrahedral subunits in a single icosahedral Pt particle. Credit: Hong Yang, University of Illinois

(Phys.org) —University of Illinois researchers have developed unique approach for the synthesis of highly uniform icosahedral nanoparticles made of platinum. Results showed that the key factors for the shape control include fast nucleation, kinetically controlled growth, and protection from oxidation by air.

"We have developed unique approach for the synthesis of highly uniform icosahedral nanoparticles made of platinum (Pt)," explained Hong Yang, a professor of chemical and biomolecular engineering at the University of Illinois at Urbana-Champaign. "This is important both in fundamental studies—nanoscience and nanotechnology—and in applied sciences such as high performance fuel cell catalysts."


Yang's research group focuses on the synthesis and understanding structure-property relationship of [nanostructured materials](#) for applications in energy, [catalysis](#), and biotechnology. Its paper, "Highly Uniform Platinum Icosahedra Made by the Hot Injection-Assisted GRAILS Method," was published this week in *Nano Letters*.



"Although polyhedral [nanostructures](#), such as a cube, tetrahedron, octahedron, cuboctahedron, and even icosahedron, have been

synthesized for several [noble metals](#), uniform Pt icosahedra do not form readily and are rarely made," stated Wei Zhou, a visiting scholar with Yang's research group and the paper's first author.

An icosahedron crystal is a [polyhedron](#) with 20 identical equilateral triangular faces, 30 edges and 12 vertices. According to Yang, icosahedral shaped crystals can improve the [catalytic activity](#) in oxygen reduction reaction partly because of the surface strain.



This is a micrograph showing the uniformity of the nanocrystals at low magnification. Credit: Hong Yang, University of Illinois

"The key reaction step to improve the activity of oxygen electrode catalysts in the [hydrogen fuel cell](#) is to optimize the bond strength between Pt and absorbed oxygen-containing intermediate species," Yang said. "This allows the rapid production of water and let the intermediate react and leave the surface quickly so the catalyst site can be used again."

"Unlike many other forms of metal nanoparticles, an icosahedral nanocrystal is not a single crystal, but has many twin (defect) boundaries within this shape. Previous simulation data suggest that it is unstable for Pt nanoparticles to exist in this shape at about >1-2 nm and, indeed, it is uncommon for Pt nanoparticles to have this morphology." Highly uniform Pt icosahedral nanocrystals with an edge length of 8.8 nm were synthesized by Yang's research group. They were made from platinum acetylacetonate in dodecylamine and with small amount of oleic acid using a hot injection-assisted GRAILS (gas reducing agent in liquid solution) approach. In the GRAILS approach, the inclusion of CO gas greatly facilitates the formation of well-defined shapes.

"Our results showed that the key factors for the shape control include fast nucleation, kinetically controlled growth, and protection from oxidation by air," Zhou added. By adjusting these key parameters, Pt hyper-branched rods, cubes, and octapods were also obtained.

"We are currently studying why this shape is formed in our systems and how we can use this principle to produce other unusual and potentially useful Pt and its alloy nanoparticles," Yang noted. "The high purity (>95%) of the products provides the ideal model materials for studying the structure/morphology-property relationships. Such mechanistic understanding is valuable for the design of advanced, high performance metal and metal alloy catalysts."

Provided by University of Illinois at Urbana-Champaign

Citation: Researchers develop unique method for creating uniform nanoparticles (2013, May 6)  
retrieved 26 April 2024 from

<https://phys.org/news/2013-05-unique-method-uniform-nanoparticles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.