


El Niño unusually active in the late 20th century, study finds

June 30 2013



This graph shows El Niño variability derived from tree rings (blue) and instrumental measurements (red). The dashed lines indicate boundary for natural variability. Recent El Niño behavior is largely beyond natural variability. Credit: International Pacific Research Center

Spawning droughts, floods, and other weather disturbances world-wide, the El Niño – Southern Oscillation (ENSO) impacts the daily life of millions of people. During El Niño, Atlantic hurricane activity wanes and rainfall in Hawaii decreases while Pacific winter storms shift southward, elevating the risk of floods in California.

The ability to forecast how ENSO will respond to [global warming](#) thus matters greatly to society. Providing accurate predictions, though, is challenging because ENSO varies naturally over decades and centuries. Instrumental records are too short to determine whether any changes

seen recently are simply natural or attributable to man-made greenhouse gases. Reconstructions of ENSO behavior are usually missing adequate records for the tropics where ENSO develops.

Help is now underway in the form of a tree-ring record reflecting ENSO activity over the past seven centuries. Tree-rings have been shown to be very good proxies for temperature and rainfall measurements. An international team of scientists spearheaded by Jinbao Li and Shang-Ping Xie, while working at the International Pacific Research Center, University of Hawaii at Manoa, has compiled 2,222 tree-ring chronologies of the past seven centuries from both the tropics and mid-latitudes in both hemispheres. Their work is published in the June 30, 2013 online issue of *Nature Climate Change*.

The inclusion of tropical tree-ring records enabled the team to generate an archive of ENSO activity of unprecedented accuracy, as attested by the close correspondence with records from equatorial Pacific corals and with an independent Northern Hemisphere temperature reconstruction that captures well-known teleconnection [climate patterns](#).



Ancient trees, such as *Polylepis tarapacana* growing in rocky soils in the South American Altiplano, are sensitive to climate anomalies associated with large-scale climate patterns stemming from the El Niño-Southern Oscillation. Reading the rings of thousands such trees lining the Pacific rim, Li and Xie found that recent El Niño activity is at its highest for the past 700 years, possibly a response to ongoing global warming. Credit: Duncan Christie

These proxy records all indicate that ENSO was unusually active in the late 20th century compared to the past seven centuries, implying that this climate phenomenon is responding to ongoing global warming.

"In the year after a large tropical volcanic eruption, our record shows that the east-central tropical Pacific is unusually cool, followed by unusual warming one year later. Like greenhouse gases, volcanic

aerosols perturb the Earth's radiation balance. This supports the idea that the unusually high ENSO activity in the late 20th century is a footprint of global warming" explains lead author Jinbao Li.

"Many climate models do not reflect the strong ENSO response to global warming that we found," says co-author Shang-Ping Xie, meteorology professor at the International Pacific Research Center, University of Hawaii at Manoa and Roger Revelle Professor at Scripps Institution of Oceanography, University of California at San Diego. "This suggests that many models underestimate the sensitivity to radiative perturbations in [greenhouse gases](#). Our results now provide a guide to improve the accuracy of climate models and their projections of future ENSO activity. If this trend of increasing ENSO activity continues, we expect to see more weather extremes such as floods and droughts."

More information: Li, J., S.-P. Xie, E. R. Cook, M. Morales, D. Christie, N. Johnson, F. Chen, R. D'Arrigo, A. Fowler, X. Gou, and K. Fang (2013): El Niño modulations over the past seven centuries. *Nature Climate Change*. [dx.doi.org/10.1038/nclimate1936](https://doi.org/10.1038/nclimate1936)

Provided by University of Hawaii at Manoa

Citation: El Nino unusually active in the late 20th century, study finds (2013, June 30) retrieved 25 April 2024 from <https://phys.org/news/2013-06-el-nino-unusually-late-20th.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.