

# New 'embryonic' subduction zone found

June 17 2013

---



Continental Europe and America could be joined in 220 million years. Image: Google maps.

(Phys.org) —A new subduction zone forming off the coast of Portugal heralds the beginning of a cycle that will see the Atlantic Ocean close as continental Europe moves closer to America.

Published in *Geology*, new research led by Monash University geologists has detected the first evidence that a passive margin in the Atlantic ocean is becoming active. Subduction zones, such as the one beginning near Iberia, are areas where one of the [tectonic plates](#) that cover the Earth's surface dives beneath another plate into the mantle - the layer just below the crust.

Lead author Dr João Duarte, from the School of [Geosciences](#) said the team mapped the [ocean floor](#) and found it was beginning to fracture, indicating tectonic activity around the apparently passive South West

Iberia plate margin.

"What we have detected is the very beginnings of an active margin - it's like an embryonic [subduction zone](#)," Dr Duarte said.

"Significant [earthquake activity](#), including the 1755 quake which devastated Lisbon, indicated that there might be convergent tectonic movement in the area. For the first time, we have been able to provide not only evidences that this is indeed the case, but also a consistent driving mechanism."

The incipient subduction in the Iberian zone could signal the start of a new phase of the Wilson Cycle - where plate movements break up supercontinents, like Pangaea, and open oceans, stabilise and then form new subduction zones which close the oceans and bring the scattered continents back together.

This break-up and reformation of supercontinents has happened at least three times, over more than four billion years, on Earth. The Iberian subduction will gradually pull Iberia towards the United States over approximately 220 million years.

The findings provide a unique opportunity to observe a passive margin becoming active - a process that will take around 20 million years. Even at this early phase the site will yield data that is crucial to refining the geodynamic models.

"Understanding these processes will certainly provide new insights on how subduction zones may have initiated in the past and how oceans start to close," Dr Duarte said.

**More information:** [geology.gsapubs.org/content/ea ...  
4100.1.full.pdf+html](http://geology.gsapubs.org/content/ea/4100.1.full.pdf+html)

Provided by Monash University

Citation: New 'embryonic' subduction zone found (2013, June 17) retrieved 23 April 2024 from <https://phys.org/news/2013-06-embryonic-subduction-zone.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.