

Researchers determine factors that influence spinach contamination pre-harvest

June 20 2013

A team of researchers from Texas and Colorado has identified a variety of factors that influence the likelihood of *E. coli* contamination of spinach on farms prior to harvest. Their research is published in the July 2013 issue of the journal *Applied and Environmental Microbiology*.

"<u>Microbial contamination</u> of produce seems strongly influenced by the time since the last <u>irrigation</u>, the workers' <u>personal hygiene</u> and the field's use prior to planting of produce," says first author Sangshin Park of Texas A&M University, College Station. "These factors, together with the role of weather in produce contamination should be the targets of future research efforts to design cost-effective strategies for control of produce contamination."

E. coli contamination of <u>spinach</u> on farms in Colorado and Texas was 172 times more likely if the produce field was within 10 miles of a poultry farm, and 64 times more likely if irrigated by pond water, says Park.

As *E. coli* is commonly used as an indicator of fecal contamination with food-borne pathogens, the practice of hygiene—availability of portable toilets and hand-washing stations for workers in the fields —and the absence of grazing or hay production on the fields prior to planting spinach, reduced the risk seven-fold.

Other potential risk factors tested in the study included numbers of workers, farm size, organic vs. conventional production, the use of



chemical fertilizers, compost, and manure, says Park. The researchers assayed 955 spinach samples from 12 farms in the two states, finding that generic *E. coli* was present on 63 of them (6.6 percent).

Of particular note, the researchers tested their statistical model for spinach contamination to determine how accurately it was able to pinpoint the level of contamination. "The assessment of the predictive performance of a developed statistical model is largely omitted from food safety studies," says Park. Their methodology may serve as a useful template for future investigations of contamination on farms, he says.

"Because produce is commonly consumed raw, it would be best to prevent pre-harvest contamination by food-borne pathogens all together or at least to reduce it," says Park.

More information: <u>www.asm.org/images/Communicati ...</u> 2013/0613spinach.pdf

Provided by American Society for Microbiology

Citation: Researchers determine factors that influence spinach contamination pre-harvest (2013, June 20) retrieved 23 May 2024 from <u>https://phys.org/news/2013-06-factors-spinach-contamination-pre-harvest.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.