

## Fishing for chips

June 25 2013

Traditional methods of marking larger farm animals rely on branding with hot irons or on ear-tagging. Dogs and cats are instead identified by the implant of a microchip transponder. With very few exceptions, it is now mandatory within the European Union to mark horses by means of transponders.

Nevertheless, some sport-horse registries oppose the use of microchips because they believe that the rate of identification failure is unacceptably high. To date, no systematic examinations to see whether chips are easy to decode, have been conducted. Manuela Wulf in the group of Christine Aurich at the Vetmeduni Vienna has therefore examined the readability of microchips in more than 400 horses. She tested each of the chips with three different scanners. The scanners differed in diameter and field strength. The scientists tested both sides of the animals' necks.

## Only high quality scanners are recommended

The results were intriguing. The "best" scanner (equipped with a <u>digital</u> <u>signal processing</u> function that filters interfering signals) detected and read all chips correctly when it was placed on the side of the neck where the <u>chip</u> was implanted and located nearly 90% of the chips even when it was on the other side of the neck. However, the other two scanners performed considerably less well, producing correct reads in around 90% of the cases when on the same side of the neck. On the opposite side of the neck, the success rate ranged between 20-25%. As Wulf puts it, "It is important that the scanners find and read the chips correctly in



every case. We can only recommend the top-of-the-range scanner, which should ideally be placed on the side of the horse's neck where the chip was implanted." However, Aurich adds, "Even the lowest quality scanner we tested, performed much better than traditional branding methods of horse identification."

## Chipping causes less injury compared to branding

The major objection to the use of branding relates to the pain and long-term damage it inflicts on the animals. Wulf and her colleagues thus investigated whether the use of microchip markers was any better. She looked closely at the site of chip implantation in 16 horses of nine different breeds and of various ages that had been submitted to the Vetmeduni for post mortem examinations. In the vast majority of cases, the chips seemed to have caused absolutely no ill effects: the two animals that were moderately affected had probably only been chipped recently and there had not yet been time for the wound to heal. As Aurich sums up the findings, "Not only is chipping a far more reliable method for marking horses than traditional methods of branding, we also found that it causes far less injury to the animals."

Provided by University of Veterinary Medicine -- Vienna

Citation: Fishing for chips (2013, June 25) retrieved 25 April 2024 from <a href="https://phys.org/news/2013-06-fishing-chips.html">https://phys.org/news/2013-06-fishing-chips.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.