

Man-made particles affect hurricane frequency, study finds

June 23 2013



A television reporter watches waves hit a pier before the arrival of Hurricane Sandy October 29, 2012 just off the Boardwalk in Atlantic City, New Jersey. Higher levels of air pollution reduced the frequency of North Atlantic hurricanes and other tropical storms for most of the 20th century, a study said Sunday.

Higher levels of air pollution reduced the frequency of North Atlantic hurricanes and other tropical storms for most of the 20th century, a study said Sunday.



Adding to evidence for mankind's impact on the <u>weather system</u>, the probe found a link between these powerful storms and aerosols, the scientific term for specks of matter suspended in a gas.

Aerosols can occur in natural form—as dusty volcanic plumes, clouds or fog—but are also man-made, such as sooty particles from burning coal or oil.

The study focused on particles from North America and Europe that were generated mainly from <u>burning fossil fuels</u>.

Researchers from the UK Met Office created weather simulations covering the period 1860 to 2050.

They found that tropical storms were much less frequent during periods when emissions of man-made aerosols increased over the North Atlantic.

"Increases in anthropogenic emissions (particularly of aerosols) through most of the last century is found to have reduced hurricane activity," coauthor Ben Booth told AFP.

"The cooling impact of man-emitted aerosols may have had a more important regional impact on climate than we previous appreciated."

Aerosols reflect solar rays and change the brightness of clouds, which affects how much of the Sun's heat is projected onto the surface of the sea, the authors suggest.

Ocean warmth provides the raw energy for tropical storms, which in <u>extreme conditions</u> can brew into destructive hurricanes.

Conversely, the study found that measures since the 1980s to tackle pollution and improve air quality reduced levels of aerosols—and in turn



ramped up hurricane activity.

"The clean-up of industrial aerosols in the last 20 years, while being beneficial for human health and linked to a recovery of African Sahel rains since the 1980s <u>droughts</u>, may have contributed to increases in Atlantic hurricane activity," Booth said by email.

The authors said their study, published in the journal *Nature Geoscience*, is the first to demonstrate a link between aerosols and Atlantic tropical storms.

The research team postulates that in the future, it will be Earth-warming greenhouse gases, much longer-lasting than aerosols, that will exert the most influence on tropical storm frequency.

Previous work published in *Nature Climate Change* had said that while the number of <u>tropical storms</u> was not projected to increase in future, their intensity was.

The hurricane season runs from June to November. For 2013, the US National Oceanic and Atmospheric Administration (NOAA) has predicted 13 to 20 "named" storms, seven to 11 hurricanes and three to six major hurricanes.

More information: Paper: <u>dx.doi.org/10.1038/ngeo1854</u> News&Views: <u>dx.doi.org/10.1038/ngeo1871</u>

© 2013 AFP

Citation: Man-made particles affect hurricane frequency, study finds (2013, June 23) retrieved 2 May 2024 from <u>https://phys.org/news/2013-06-man-made-particles-affect-hurricane-frequency.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.