


Resistivity switch is window to role of magnetism in iron-based superconductors

June 5 2013



A polarized light microscopy image of structural domains in a barium-iron-arsenide crystal, a type of superconductor. Physicists at the US Department of Energy's Ames Laboratory discovered unexpected resistivity changes in crystals doped with potassium, which suggest a close relationship between magnetism and superconductivity in iron-based superconductors. Credit: U.S. Dept. of

(Phys.org) —Physicists at the U.S. Department of Energy's Ames Laboratory have discovered surprising changes in electrical resistivity in iron-based superconductors. The findings, reported in *Nature Communications*, offer further evidence that magnetism and superconductivity are closely related in this class of novel superconductors.

"We found that the directions of smallest and largest resistivity within the conducting layers are significantly dependent on the composition of the compounds, and in some compositions, they change sign, or, in other words, electric current flows easier in the direction that was originally more difficult" said Ames Laboratory faculty scientist Ruslan Prozorov. "This change can only be explained if the underlying [magnetic behavior](#) is intimately connected to superconductivity."


Understanding the basic physics behind iron-based superconductivity may someday make it possible to use them for super-efficient energy technologies.

Erick Blomberg, an Ames Laboratory graduate research assistant, spearheaded the research under the guidance of Prozorov and Ames Laboratory scientist Makariy Tanatar. The team studied resistivity in [barium](#)-iron-arsenide compounds with different chemical substitutions, which allowed for probing [electronic properties](#) throughout the entire regime in which magnetism and superconductivity coexist.

In [conventional superconductors](#), magnetism and superconductivity do not coexist and one suppresses another. In contrast, some iron-based superconductors show significant overlap between magnetism and

superconductivity, which allows for unique measurements of their properties in the coexistence region.

"But there is a complication," said Prozorov, who is also an Iowa State University professor in the Department of Physics & Astronomy. "In the coexistence region, the crystal structure of barium-iron-arsenide crystals is such that the crystal splits into the structural domains (also known as structural twins), which makes studying directional electronic properties difficult.



(left to right) Erick Blomberg, Makariy Tanatar and Ruslan Prozorov.

To eliminate the structural domains, the team developed a technique in which single crystals are physically pulled apart. A sample is suspended by wires in a horseshoe-shaped bracket, which can be mechanically

stretched. This assembly is then placed into a small liquid helium cryostat to reach temperatures where magnetism and [superconductivity](#) coexist. Polarized optical microscopy is used to distinguish between different structural domains to verify that the samples are in the detwinned (without fragmented domains) state. Electric resistivity is then measured.

Blomberg and colleagues studied a series of samples, covering a precise range of compositions, provided by Nanjing University's Hai-Hu Wen.

"In this case, making chemical substitutions was, in effect, taking electrons away from the compound or adding electrons to the compound," said Blomberg, who will soon receive a PhD in condensed matter physics from Iowa State University. "And as you take electrons away, at a certain point the direction in which it was harder for electricity to flow becomes the direction it was easier for it to flow."

Theoretical predictions made by physicists Jörg Schmalian and Rafael Fernandez (both former Ames Laboratory scientists) along with Naval Research Lab's Igor Mazin and Michelle Johannes, closely matched the experimental findings.

More information: [www.nature.com/ncomms/journal/ ... full/ncomms2933.html](http://www.nature.com/ncomms/journal/full/ncomms2933.html)

Provided by Ames Laboratory

Citation: Resistivity switch is window to role of magnetism in iron-based superconductors (2013, June 5) retrieved 20 March 2024 from <https://phys.org/news/2013-06-resistivity-window-role-magnetism-iron-based.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.