

# Scientists say Russian meteor shockwave went twice around globe

June 29 2013, by Nancy Owano

---



Meteor trail over eastern Russia is seen in this image from the Russian Emergency Ministry. Credit: Russian Emergency Ministry

(Phys.org) —Scientists have determined that an [asteroid](#) that burned up over Russia in February resulted in a shock wave so powerful that it traveled twice around the globe. They made that determination by means of a system of sensors used to detect evidence of nuclear tests. Researchers studied data from the International Monitoring System

network operated by the Comprehensive Nuclear-Test-Ban Treaty Organization. Its stations monitor ultra-low frequency acoustic waves, or infrasound, from nuclear test explosions. The system can also detect blasts from other sources. According to BBC News, it was the most powerful event ever recorded by the network.

The 10,000-ton space rock burned up above Chelyabinsk. The BBC said Alexis Le Pichon, Atomic Energy Commission in France, and his research colleagues reported that the explosive energy of the impact was equivalent to 460 kilotons of TNT. Writing in the *Geophysical Research Letters* journal, their article is titled "Russian Fireball Largest Ever Detected by CTBTO infrasound sensors." Authors are Alexis Le Pichon, Lars Ceranna, Christoph Pilger, Pierrick Mialle, David Brown, Pascal Herry, and Nicolas Brachet.

"On 15 February 2013," they wrote, "a large Earth-impacting fireball disintegrated over the Ural Mountains. This extraordinary event is, together with the 1908 Tunguska fireball, among the most energetic events ever instrumentally recorded." The 1908 reference is to the event that occurred that year in Siberia.

As for the February event, the authors noted that, after circling the globe, it generated infrasound returns at distances up to ~85,000 km, and was detected at 20 infrasonic stations of the global International Monitoring System. Multiple arrivals involving waves that traveled twice round the globe were identified.

"A preliminary estimate of the explosive energy using empirical period-yield scaling relations gives a value of 460 kilotons of TNT equivalent. In the context of the future verification of the Comprehensive Nuclear-Test-Ban Treaty (CTBT), this event provides a prominent milestone for studying in detail infrasound propagation around the globe for almost three days as well as for calibrating the performance of the IMS

network."

Their numbers suggest the February [asteroid](#) event is the most energetic event reported since the 1908 Tunguska meteor in Siberia. Turning to Tunguska, another team of scientists published a study offering their evidence of the meteoritic origin of the Tunguska cosmic body. Their discussion of the 1908 fireball is in the journal *Planetary and Space Science*. Diamond–lonsdaleite–graphite micro-samples collected from peat after the 1908 catastrophic blast in the Tunguska area were studied with scanning (SEM) and transmission electron (TEM) microscopy, NanoSecondary Ion Mass Spectrometry (NanoSIMS) and an X-ray synchrotron technique.

**More information:** *Geophysical Research Letters* [DOI: 10.1002/grl.50619](#)  
*Planetary and Space Science* [DOI: 10.1016/j.pss.2013.05.003](#)

© 2013 Phys.org

Citation: Scientists say Russian meteor shockwave went twice around globe (2013, June 29) retrieved 27 March 2023 from <https://phys.org/news/2013-06-scientists-russian-meteor-shockwave-globe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--