

Capturing black hole spin could further understanding of galaxy growth

July 29 2013



This is a j-peg artist's impression of a supermassive black hole at the center surrounded by matter flowing onto the black hole in what is termed an accretion disk. Also shown is an outflowing jet of energetic particles, believed to be powered by the black hole's spin. Credit: NASA/JPL-Caltech

Astronomers have found a new way of measuring the spin in supermassive black holes, which could lead to better understanding about how they drive the growth of galaxies.

The scientists at Durham University, UK, observed a black hole - with mass 10 million times that of our Sun - at the centre of a [spiral galaxy](#)

500 million light years from Earth while it was feeding on the surrounding disc of material that fuels its growth and powers its activity.

By viewing optical, ultra-violet and soft [x-rays](#) generated by heat as the black hole fed, they were able to measure how far the disc was from the black hole.

This distance depends on black hole spin as a fast spinning black hole pulls the disc in closer to itself, the researchers said. Using the distance between the black hole and the disc, the scientists were able to estimate the spin of the black hole.

The scientists said that understanding spin could lead to greater understanding of galaxy growth over billions of years.

Black holes lie at the centres of almost all [galaxies](#), and can spit out incredibly hot particles at high energies that prevent intergalactic gases from cooling and forming [new stars](#) in the outer galaxy. Scientists don't yet understand why the jets are ejected into space, but the Durham experts believe that their power could be linked to the spin of the black hole.

Spin is difficult to measure as it only affects the behaviour of material really close to the black hole.

The research, funded by the Science and Technology Facilities Council, is published in *Monthly Notices of the Royal Astronomical Society*.

Lead researcher Professor Chris Done, in the Department of Physics, at Durham University, said: "We know the black hole in the centre of each galaxy is linked to the galaxy as a whole, which is strange because black holes are tiny in relation to the size of a galaxy. This would be like something the size of a large boulder (10m), influencing something the

size of the Earth.

"Understanding this connection between stars in a galaxy and the growth of a black hole, and vice-versa, is key to understanding how galaxies form throughout cosmic time.

"If a black hole is spinning it drags space and time with it and that drags the accretion disc, containing the black hole's food, closer towards it. This makes the black hole spin faster, a bit like an ice skater doing a pirouette.

"By being able to measure the distance between the black hole and the accretion disc, we believe we can more effectively measure the spin of black holes.

"Because of this, we hope to be able to understand more about the link between [black holes](#) and their galaxies."

The Durham scientists were able to measure the spin of the black hole using soft x-ray, optical and ultra-violet images captured by the European Space Agency's XMM-Newton satellite.

They said further research was needed to reduce the effects of black hole mass on their findings.

More information: A new way to measure supermassive black hole spin in accretion disc dominated Active Galaxies, Done C, et al, *Monthly Notices of the Royal Astronomical Society*, [dx.doi.org/10.1093/mnras/stt1138](https://doi.org/10.1093/mnras/stt1138)

Provided by Durham University

Citation: Capturing black hole spin could further understanding of galaxy growth (2013, July 29)
retrieved 23 April 2024 from

<https://phys.org/news/2013-07-capturing-black-hole-galaxy-growth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.