

New knowledge about permafrost improving climate models

July 28 2013

New research findings from the Centre for Permafrost (CENPERM) at the Department of Geosciences and Natural Resource Management, University of Copenhagen, document that permafrost during thawing may result in a substantial release of carbon dioxide into the atmosphere and that the future water content in the soil is crucial to predict the effect of permafrost thawing. The findings may lead to more accurate climate models in the future.

The permafrost is thawing and thus contributes to the release of carbon dioxide and other [greenhouse gases](#) into the atmosphere. But the rate at which carbon dioxide is released from permafrost is poorly documented and is one of the most important uncertainties of the current [climate](#) models.

The knowledge available so far has primarily been based on measurements of the release of carbon dioxide in short-term studies of up to 3-4 months. The new findings are based on measurements carried out over a 12-year period. Studies with different water content have also been conducted. Professor Bo Elberling, Director of CENPERM (Centre for Permafrost) at the University of Copenhagen, is the person behind the novel research findings which are now being published in the internationally renowned scientific journal *Nature Climate Change*.

"From a [climate change](#) perspective, it makes a huge difference whether it takes 10 or 100 years to release, e.g., half the permafrost carbon pool. We have demonstrated that the supply of oxygen in connection with

drainage or drying is essential for a rapid release of carbon dioxide into the atmosphere," says Bo Elberling.

Water content in the soil crucial to predict effect of permafrost thawing

The new findings also show that the future water content in the soil is a decisive factor for being able to correctly predict the effect of permafrost thawing. If the permafrost remains water-saturated after thawing, the carbon decomposition rate will be very low, and the release of carbon dioxide will take place over several hundred years, in addition to methane that is produced in waterlogged conditions. The findings can be used directly to improve existing [climate models](#).

The new studies are mainly conducted at the Zackenberg research station in North-East Greenland, but permafrost samples from four other locations in Svalbard and in Canada have also been included and they show a surprising similarity in the loss of carbon over time.

"It is thought-provoking that microorganisms are behind the entire problem – microorganisms which break down the carbon pool and which are apparently already present in the permafrost. One of the critical decisive factors – the [water content](#) – is in the same way linked to the original high content of ice in most permafrost samples. Yes, the temperature is increasing, and the permafrost is thawing, but it is, still, the characteristics of the [permafrost](#) which determine the long-term release of [carbon dioxide](#)," Bo Elberling concludes.

More information: *Nature Climate Change* [DOI: 10.1038/nclimate1955](https://doi.org/10.1038/nclimate1955)

Provided by University of Copenhagen

Citation: New knowledge about permafrost improving climate models (2013, July 28) retrieved 9 April 2024 from <https://phys.org/news/2013-07-knowledge-permafrost-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.