


# Researchers perform first direct measurement of Van der Waals force

July 8 2013, by Bob Yirka



Mapping out the van der Waals interaction between two atoms. (a) In the experiment of Béguin et al. two atoms are trapped in the foci of two laser beams separated by a distance  $R$ . (b) Depending on  $R$ , the excitation laser field can couple the ground state  $|gg\rangle$  of the atomic pair to states containing one atom in the Rydberg state ( $|rg\rangle$  and  $|gr\rangle$ , respectively), or to a state with both atoms populating the Rydberg state  $|rr\rangle$ . The energy of the latter state is strongly shifted because of the van der Waals interaction  $U_{\text{vdW}}$  between the atoms (see level diagram in the upper left), resulting in a distance-dependent coherent excitation dynamics of the atomic pair system. (c) By analyzing the time evolution of the atom-pair state, Béguin et al. deduce the van der Waals energy shift as a function of interatomic distance for different Rydberg states. Credit: *Physics* 6, 71 (2013) DOI: 10.1103/Physics.6.71

(Phys.org) —Researchers working at the French National Center for Scientific Research have for the first time, directly measured the Van der Waals force between two atoms. In their paper published in the journal *Physical Review Letters*, the team describes how they used lasers to hold two atoms steady and a third laser to measure the Van der Waals

force between them.

The weak force between atoms, named after its discoverer Johannes Diderik van der Waals is evident in the behavior of many materials—it's what keeps most [gas molecules](#) together. Scientists have also discovered that it's also what allows a gecko's toes to stick to a smooth wall. But, because the [weak force](#) is only apparent when atoms are very close together, scientists have until now been unable to measure it directly.

In this new effort, the research team chose to use Rydberg atoms as part of their study. Such atoms are large and one of their electrons has a highly charged state. This makes them a good candidate for attempting to measure the Van der Waals force—they have more force between them than most other atom pairs and because of that can be measured at longer distances.

They started by firing a pair of lasers at twin Rydberg atoms. Doing so held them steady. Next, they fired a third laser at the two atoms causing them to oscillate at a desired frequency. By measuring the oscillations, the researchers were able to work out mathematically the Van der Waals force between them. More specifically, the researchers measured [oscillations](#) between ground and excited states, noting that the distance between the two atoms at the time of measurement was key—too close and the excitation of one of the atoms overwhelmed the other—too far and the force between the atoms became too weak to measure. Using the third laser as optic tweezers allowed for adjusting the distance between the two atoms, which ultimately led to just the right distance for measurement.

The team also noted that the technique used for measuring the Van der Waals force also led to the oscillating atoms evolving to a fully coherent state. This means the technique could be used to create quantum logic gates, which might prove useful in creating a quantum computer.

**More information:** Direct Measurement of the van der Waals Interaction between Two Rydberg Atoms, *Phys. Rev. Lett.* 110, 263201 (2013) [prl.aps.org/abstract/PRL/v110/i26/e263201](http://prl.aps.org/abstract/PRL/v110/i26/e263201)

### **Abstract**

We report the direct measurement of the van der Waals interaction between two isolated, single Rydberg atoms separated by a controlled distance of a few micrometers. Working in a regime where the single-atom Rabi frequency for excitation to the Rydberg state is comparable to the interaction, we observe partial Rydberg blockade, whereby the time-dependent populations of the various two-atom states exhibit coherent oscillations with several frequencies. Quantitative comparison of the data with a simple model based on the optical Bloch equations allows us to extract the van der Waals energy, and observe its characteristic  $C_6/R^6$  dependence. The measured  $C_6$  coefficients agree well with ab initio calculations, and we observe their dramatic increase with the principal quantum number  $n$  of the Rydberg state.

© 2013 Phys.org

Citation: Researchers perform first direct measurement of Van der Waals force (2013, July 8) retrieved 24 April 2024 from <https://phys.org/news/2013-07-van-der-waals.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--