


New risk model sheds light on arsenic risk in China's groundwater

August 22 2013

Modeled probability of groundwater arsenic contamination in China exceeding the WHO guideline for drinking water of 10 µg/L




This is modeled probability of groundwater arsenic contamination in china exceeding the WHO guideline for drinking water of 10 microgramms/liter. Credit: Rodriguez-Lado et al.; Science 341 (6148)

Arsenic-laden groundwater used for cooking and drinking could pose a risk to the health of almost 20 million people across China. This is shown by a study carried out by Eawag scientists in collaboration with

Chinese colleagues and published today in *Science*. The estimates are based on a risk model incorporating geological and hydrological data, as well as measurements of arsenic in wells. The study is being adopted by the authorities in the national groundwater monitoring program.

Since the 1960s, it has been known that [groundwater](#) resources in certain provinces of China are contaminated with arsenic. Estimates of the numbers of affected people have risen year by year. In the most recent survey – conducted by the Chinese Ministry of Health between 2001 and 2005 – more than 20,000 (5%) of the 445,000 wells tested showed arsenic concentrations higher than 50 µg/L. According to official estimates, almost 6 million people consume drinking water with an arsenic content of more than 50 µg/L and almost 15 million are exposed to concentrations exceeding 10 µg/L (the guideline value recommended by the WHO).

Given the sheer size of China and the time and expense involved in testing for arsenic contamination, several more decades would probably be required to screen all of the millions of groundwater wells. Accordingly, a group of researchers from Eawag and the China Medical University in Shenyang de-veloped a statistical risk model making use of existing data on geology, soil characteristics and topographic features. This model was calibrated using available arsenic measurements. The predictions of unsafe or safe areas showed a high level of agreement, both for known high-risk areas and for areas where elevated [arsenic levels](#) had been ruled out by sampling campaigns.



This shows the estimated population at potential risk of arsenic exposure over 10 microgramms/liter from groundwater-derived drinking water in China. Credit: Rodriguez-Lado et al.; Science 341 (6148)

In addition, large areas have now been identified as potentially at risk, such as the basins of the Tarim (Xinjiang), Ejina (Inner Mongolia) and Heihe (Gansu), or the North China Plain (Henan and Shandong). Arsenic concentrations above 10 µg/L are predicted for a total area of 580,000 km². When these results were combined with the latest available population data, it was found that almost 20 million people across China live in high-risk areas. Geochemist Annette Johnson concedes: "This figure may be an overestimate, as we lack reliable information on the number of people with treated water supplies." But in the long term, she adds, China will remain dependent on groundwater as a source of drinking water, particularly in the arid provinces. The [risk model](#) shows where conventional groundwater quality monitoring efforts are best

focussed: "Our method permits more targeted sampling campaigns and saves time in identifying populations at risk. The Chinese authorities are adopting our maps in the national monitoring programme." Johnson is convinced that the model could also be used in other countries where groundwater is known or suspected to be contaminated with arsenic – for example, in Africa or in central Asia, where risk assessments for [arsenic contamination](#) have not yet been performed.

Arsenic

Arsenic is one of the most common inorganic contaminants found in drinking water world-wide. This metalloid occurs as a natural component of sediments, with small quantities being dissolved in groundwater as a result of weathering. The inorganic salts of arsenic are tasteless and odourless, but highly toxic to humans. If ingested over long periods, even low concentrations can cause damage to health, including hyperpigmentation of the skin, hyperkeratosis on the palms and soles, disorders of liver, cardiovascular and kidney function, and various types of cancer. Problems arise from the fact that firstly, arsenic concentrations can vary widely at the local level and, secondly, people in many areas are completely unaware of the risk because their groundwater wells have never been screened for arsenic. Concentrations below 10 µg/L are considered safe. This concentration is therefore recommended by the World Health Organization as a guideline value for arsenic in [drinking water](#). In China, the standard guideline has just recently changed from 50 µg/L to 10 µg/L. In many other studies, e.g. at various sites in Inner Mongolia, [arsenic](#) concentrations of more than 100 µg/L and up to 1500 µg/L have been measured.

Provided by Swiss Federal Institute of Aquatic Science and Technology:
Eawag

Citation: New risk model sheds light on arsenic risk in China's groundwater (2013, August 22)
retrieved 26 April 2024 from <https://phys.org/news/2013-08-arsenic-china-groundwater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.