

Honeyguide birds destroy own species' eggs to eliminate competition

August 20 2013



This is a bee-eater clutch parasitised by a honeyguide; the larger egg is a honeyguide egg, and the smaller eggs are all rotten because they have been punctured by the laying female honeyguides. Credit: Claire Spottiswoode, University of Cambridge

Like cuckoos, honeyguides are parasitic birds that lay their eggs in other birds' nests and dupe them into raising their young. Now scientists reveal that, unlike in cuckoos, the resemblance between honeyguide eggs and

those of their bee-eater bird hosts hasn't evolved to trick hosts into accepting the imposter egg as one of their own. Rather, it appears to have evolved to trick other honeyguides who would otherwise destroy the eggs because of fierce competition for host nests. The new research is published today, 21 August, in the journal *Biology Letters*.

Honeyguides are intriguingly odd birds that are best-known for their unique, mutually [beneficial relationship](#) with humans. Honeyguides love to eat beeswax. To obtain it, they guide human honey-hunters to bees' nests. In return for showing the humans the bees, the honeyguide gains access to the otherwise dangerous and impenetrable nest and its sought-after wax.

But these African birds also have a dark side. They are unusually vicious [parasites](#) whose imposter chicks stab the chicks of their host birds (often little bee-eaters) to death as soon as they hatch in order to eliminate competition for the host parents' care. The newly published research has shown that this fight for monopoly of the nest also extends to other honeyguides in a battle conducted deep underground in the nest burrows that bee-eaters dig into the roofs of Aardvark holes.

The researchers' curiosity was piqued by their earlier finding that like cuckoo eggs, honeyguide eggs resemble those of each of their several [host species](#). Instead of mimicking their colour, however, they mimic their size (as colour is irrelevant in the dark interior of the deep holes in which hosts breed). For example, honeyguides parasitising little bee-eaters lay smaller eggs in their nests than do honeyguides parasitising larger hosts. Many classic studies have shown that comparable [mimicry](#) in cuckoo eggs has evolved to reduce rejection by choosy hosts that eject mismatched eggs from their nests.

Dr Claire Spottiswoode from the University of Cambridge's Department of Zoology, who carried out the research, said: "I assumed honeyguide

egg mimicry had evolved just like [cuckoo](#) mimicry, so was bowled over and baffled when little bee-eaters turned out to be pretty dim. When I played the honeyguide and experimentally parasitised their nests, the bee-eaters blithely incubated eggs even much larger than their own. So I was quite wrong, and mimicry probably hasn't evolved to dupe bee-eaters.

"It was only when some of my experiments were parasitised by real honeyguides, and my experimental eggs were pecked to pieces, that the penny dropped – perhaps they need to look like bee-eaters or else other honeyguides will get rid of them, to avoid suffering the same grisly fate that they impose upon their hosts."

A second experiment supported this hypothesis: again the researchers placed a larger foreign egg in bee-eaters nests, but then waited for a real honeyguide to come along and lay her own egg. When this happened they counted the number of puncture holes the female honeyguide made in each egg in the clutch. This revealed that laying honeyguides punctured the larger foreign egg more heavily than host eggs, lest it be another honeyguide's egg and kill their own chick should it hatch first. Evolution should then favour honeyguide females that lay eggs resembling those of bee-eaters and thereby avoid being detected and destroyed by a competing honeyguide.

The suggestion that mimicry might evolve because parasites benefit from concealing their eggs from one another was first made by Cambridge scientists Nick Davies and Michael Brooke 25 years ago, but this study is the first to show that it probably happens in the wild. Honeyguide parasitism on little bee-eaters is very common and about a third of parasitised nests contain [eggs](#) laid by two or more honeyguide females, resulting in especially strong parasitic competition.

Dr Spottiswoode said: "Under these circumstances it makes good sense that honeyguides have a lot to gain from tricking other honeyguides. But

we still don't know why bee-eaters parents themselves are so indiscriminating, especially when they pay such high costs of being conned – all your offspring hacked to death, and over a month wasted raising the wrong chick!"

The research was funded by The Royal Society and the DST/NRF Centre of Excellence at Percy FitzPatrick Institute, University of Cape Town. It forms part of a wider research programme investigating coevolution between parasitic birds and their hosts in Zambia led by Dr Claire Spottiswoode, who adds "My colleagues and I are very lucky to be helped by a wonderful team of local field assistants who find all the nests we study, and in this case dug many holes into rock-hard soil to reach bee-eater [nests](#) and carry out these experiments!"

More information: The paper 'A brood parasite selects for its own egg traits' will be published in the 21 August edition of *Biology Letters*: [rsbl.royalsocietypublishing.org ... /1098/rsbl.2013.0573](http://rsbl.royalsocietypublishing.org/.../1098/rsbl.2013.0573)

Provided by University of Cambridge

Citation: Honeyguide birds destroy own species' eggs to eliminate competition (2013, August 20) retrieved 23 April 2024 from

<https://phys.org/news/2013-08-honeyguide-birds-species-eggs-competition.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.