


Relationship between the ozone depletion and the extreme precipitation in austral summer

August 29 2013



Changes due to ozone hole of (left) mean precipitation and (right) 99th percentile daily precipitation

The new study by Prof. Sarah Kang from Ulsan National Institute of Science and Technology (UNIST), showed that the ozone depletion over the South Pole has affected the extreme daily precipitation in the austral summer, for December, January, and February (DJF). This work was published in the journal *Geophysical Research Letter*.

The ozone hole over the Antarctic has affected [atmospheric circulation](#) in the Southern Hemisphere (SH) summer and Kang's previous article in *Science*, analyzed the impact of ozone depletion to increased rainfall in the subtropics.

The new article is explaining about the impact of [stratospheric ozone depletion](#) on the extreme daily precipitation in the austral summer with two [global climate models](#): the Canadian middle Atmosphere Model (CMAM) and the National Center for Atmospheric Research Community Atmospheric Mode (CAM3).

This research study has also been highlighted in the journal *Nature Geoscience* in September.

The research team led by Prof. Kang focused on a carefully conceived set of multimodel integrations forced only with observed stratospheric ozone changes. This single-forcing approach allows the research team to show extremes, in response to stratospheric ozone depletion and that these changes are likely of a dynamic rather than thermodynamic nature.

The [ozone layer](#) is a layer in the Earth's atmosphere which absorbs most of the UV radiation and contains relatively high concentrations of ozone (O₃). This ozone layer had been broken by the widespread use of man-made compounds containing chlorofluorocarbons (CFCs). After the discovery of the ozone hole, 1989 Montreal Protocol signed by 196 countries to reduce global CFC production to protect the ozone hole

In this research they discovered that the ozone depletion in the Antarctic area is associated with extreme rain in the austral summer and it would be used to forecast heavy rain and the natural disasters in the future.

Dangerous floods have occurred in every Australian state over the last 150 years. Some caused great loss of life, others devastated infrastructure. Between 1852 and 2011 at least 951 people were killed by floods, another 1326 were injured, and the cost of damage reached an estimated \$4.76 million dollars.

Even though we can predict these natural disasters in advance, we can't

stop the flood but we can be prepared for it and reduce the damage.

"Due to limited data availability in the SH, it is hard to robustly determine observed changes in extreme precipitation," said Prof. Kang. "However, since the recent Southern Hemisphere climate change is driven by the [ozone hole](#), we can deduce the recent trend from our climate model integrations."

"We would expand our research to see the correlation of the ozone depletion of the North Pole and the climate changes in the Northern Hemisphere," said Prof. Kang, showing her future research plan.

More information: "Modeling evidence that ozone depletion has impacted extreme precipitation in the austral summer", Vol. 40, 1-6, [DOI: 10.1002/grl.50796](https://doi.org/10.1002/grl.50796) , 2013.

Provided by Ulsan National Institute of Science and Technology

Citation: Relationship between the ozone depletion and the extreme precipitation in austral summer (2013, August 29) retrieved 18 April 2024 from <https://phys.org/news/2013-08-relationship-ozone-depletion-extreme-precipitation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--