

Heat intensifies Siberian wildfires

August 7 2013



The map above shows land surface temperature anomalies for July 20, 2013. Rather than depicting absolute temperatures, the map shows how much the temperatures for that week differed from the long-term average for the area. Credit: Jeff Schmaltz, LANCE/EOSDIS MODIS Rapid Response Team at NASA GSFC

The summer of 2012 was the most severe wildfire season Russia had faced in a decade. 2013 might be headed in the same direction after an unusual heat wave brought a surge of fire activity in northern Siberia in July.



A persistent high-pressure weather pattern in the Russian Arctic—a blocking high—contributed to the heat wave, which saw temperatures reach 32° Celsius (90° Fahrenheit) in the northern city of Norilsk. For comparison, daily July highs in Norilsk average 16° Celsius (61° Fahrenheit). Blocking highs are so named because they block the jet stream from moving rain-bearing weather systems along their normal west-to-east path; this leads to "stuck" weather patterns with long periods of stable air and exceptional heat.

The map above shows land surface temperature anomalies for July 20

Provided by NASA's Goddard Space Flight Center

Citation: Heat intensifies Siberian wildfires (2013, August 7) retrieved 4 June 2024 from <u>https://phys.org/news/2013-08-siberian-wildfires.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.