

## **Controlling skyrmions for better electronics**

August 8 2013



The image shows a magnetic skyrmion in a hexagonal atom arrangement (transparant spheres). The cones represent the magnetic moments and point up (red) in the skyrmion center and down (green) around it. The rotational sense of the skyrmion for a given system is always the same, as the other rotational sense has a higher energy due to an anisotropic exchange interaction. The magnetic skyrmion in PdFe/Ir(111) that can be created and annihilated with small currents from a scanning tunneling microscope tip consists of roughly 270 surface atoms. Credit: Prof. R. Wiesendanger, University of Hamburg



Physicists at the University of Hamburg managed for the first time to individually write and delete single skyrmions, a knot-like magnetic entity. Such vortex-shaped magnetic structures exhibit unique properties which make them promising candidates for future data storage devices. Skyrmions have been in the focus of active research for the last years; however, up to now these states have been merely investigated, a controlled manipulation has not been achieved.

Now this has been realized by scientists from the group of Prof. Roland Wiesendanger in Hamburg, Germany. As *Science* reports online on August 8, 2013, the creation and annihilation of single skyrmions, corresponding to writing and deleting of information on a <u>storage</u> <u>medium</u>, has been demonstrated by these researchers. This work solves one of the longstanding technical problems concerning the future use of skyrmions in information technology.

Future electronic devices are expected to become smaller while increasing their data <u>storage capacities</u> at the same time. This will soon bring classical <u>storage technologies</u> to their physical limits. In conventional <u>memory devices</u> used up to now, magnetic bits consist of many atoms with their magnetic moments aligned parallel to each other like bar magnets. Pointing in defined directions, they can represent the values "1" and "0" which are the basis for information technology. With the continuing miniaturization, the interaction between neighboring bits increases due to magnetic stray fields which can lead to loss of data. In addition, small <u>magnetic bits</u> are less stable against <u>thermal fluctuations</u> which is also called the superparamagnetic limit.

Using more "robust" <u>magnetic structures</u> like skyrmions could be a way out of this technological dead end. These structures can be imagined as a two-dimensional knot in which the magnetic moments are rotating about 360° in a plane with a unique rotational sense (see Fig. above). These particle-like magnetic knots can be assigned a kind of charge, the



topological charge. With this it is possible to assign the bit states "1" and "0" to the existence or non-existence of a skyrmion.

A clever choice of temperature and external magnetic field enabled the scientists of the group of Prof. Roland Wiesendanger to prepare and manipulate single magnetic skyrmions for the first time. They used a two atomic layer thick film of palladium and iron on an iridium crystal. In an external magnetic field this sample exhibits single localized skyrmions with a diameter of a few nanometers that can be imaged in a spin-polarized scanning tunneling microscope. Using small currents from the tip of the microscope these skyrmions can be written and deleted. For the creation the previously parallel <u>magnetic moments</u> are twisted to form a knot-like magnetic state, and for the deletion this knot is unwound again.



Spin-resolved scanning tunneling microscopy (STM) image of single magnetic



skyrmions in a bilayer of palladium and iron on top of an iridium crystal, colorized with a difference image to the same area without skyrmions. A magnetic STM tip, which is sensitive mostly to the in-plane magnetization component of the sample as indicated, is used. The cones indicate the magnetization direction in the sample. Skyrmions can be created and annihilated locally by injecting spin-polarized currents from the STM tip. Color scale: green and red correspond to magnetic moments parallel and antiparallel to the tip magnetization, respectively. Measurement parameters: U = +250 mV, I = 1 nA, B = +3.25 T, T = 4.2 K. Credit: Prof. R. Wiesendanger, University of Hamburg

"We finally found a magnetic system in which we can locally switch between ordinary ferromagnetic order and a complex spin configuration", says Dr. Kirsten von Bergmann, senior scientist in the Wiesendanger group. As published in the current issue of the *Science* magazine, four skyrmions were specifically addressed and could be created and annihilated. "We transferred the idea of tying a knot to memorize something to the field of storage technology so we can now store data in a two-dimensional magnetic knot", explains Ph.D. student Niklas Romming.

Whether skyrmions will be used as <u>data storage</u> units in our computers, tablets, or smartphones, is not foreseeable. The experimentally accomplished writing and deleting of skyrmions, however, has demonstrated the feasibility of this technology and paved the way towards a realization of such devices.

More information: "Writing and Deleting Single Magnetic Skyrmions" *Science* 9 August 2013: Vol. 341 no. 6146 pp. 636-639 DOI: 10.1126/science.1240573



## Provided by University of Hamburg

Citation: Controlling skyrmions for better electronics (2013, August 8) retrieved 3 May 2024 from <u>https://phys.org/news/2013-08-skyrmions-electronics.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.