

Wildfires projected to worsen with climate change

August 28 2013



A fire burns in Sequoia & Kings National Park, Calif., in 2005. Research by environmental scientists at Harvard University predicts wildfire seasons by 2050 will be three weeks longer, up to twice as smoky, and will burn a wider area in the western United States. Credit: National Park Service/Parker Bevington.

Research by environmental scientists at the Harvard School of

Engineering and Applied Sciences (SEAS) brings bad news to the western United States, where firefighters are currently battling dozens of fires in at least 11 states.

The Harvard team's study suggests wildfire seasons by 2050 will be about three weeks longer, up to twice as smoky, and will burn a wider area in the western states. The findings are based on a set of internationally recognized [climate scenarios](#), decades of historical [meteorological data](#), and records of past [fire activity](#).

The results will be published in the October 2013 issue of *Atmospheric Environment* and are available in advance online.

Awareness is building that gradual [climate change](#) may contribute in the coming years to increases in significant, disruptive events like [severe storms](#) and floods. Loretta J. Mickley, a senior research fellow in [atmospheric chemistry](#) at Harvard SEAS and coauthor of the new study, is thinking one step further, to secondary effects like [forest fires](#) and air quality that rely heavily on meteorological factors.


"We weren't altogether certain what we would find when we started this project," Mickley says. "In the future atmosphere we expect warmer temperatures, which are conducive to fires, but it's not apparent what the rainfall or relative humidity will do. Warmer air can hold more water vapor, for instance, but what does this mean for fires?"

"It turns out that, for the western United States, the biggest driver for fires in the future is temperature, and that result appears robust across models," Mickley adds. "When you get a large [temperature increase](#) over time, as we are seeing, and little change in rainfall, fires will increase in size."

Reaching that conclusion with statistical confidence required months of

analysis, because at the local level, wildfires are very difficult to predict.

Percentage increase in organic carbon particles in summer, mid-21st century compared to 2000



Organic carbon is one component of smoke from wildfires. Overall, wildfires in 2050 are expected to be up to twice as smoky, threatening visibility and public health. Credit: Xu Yue.

"Wildfires are triggered by one set of influences—mainly human activity and lightning—but they grow and spread according to a completely different range of influences that are heavily dependent on the weather," says lead author Xu Yue. "Of course, when all the factors come together just right—whoosh, there's a big fire."

By examining records of past weather conditions and wildfires, the team found that the main factors influencing the spread of fires vary from region to region. In the Rocky Mountain Forest, for example, the best predictor of wildfire area in a given year is the amount of moisture in the forest floor, which depends on the temperature, rainfall, and relative humidity that season. In the Great Basin region, different factors apply. There, the area burned is influenced by the relative humidity in the previous year, which promotes fuel growth. Yue, who was a postdoctoral fellow at Harvard SEAS and is now at Yale University, created mathematical models that closely link these types of variables—seasonal temperatures, [relative humidity](#), the amount of dry fuel and so forth—with the observed wildfire outcomes for six "ecoregions" in the West.

After developing those models, the team replaced the historical observations with data based on the conclusions of the fourth Intergovernmental Panel on Climate Change (IPCC), which use socioeconomic scenarios to predict possible future atmospheric and climatological conditions. For this study, the Harvard group followed the A1B scenario, which considers the climatological effect of a fast-growing global economy relying on a mixture of fossil fuels and renewable energy sources. By running the IPCC's climate data for the year 2050 through their own fire prediction models, the Harvard team was able to calculate the area burned for each ecoregion at midcentury.

For example, the calculations suggest the following for 2050 in the western United States, in comparison to present-day conditions:


- The area burned in the month of August could increase by 65% in the Pacific Northwest, and could nearly double in the Eastern Rocky Mountains/Great Plains regions and quadruple in the Rocky Mountain Forest region.
- The probability of large fires could increase by factors of 2-3.

- The start date for the fire season could be earlier (late April instead of mid-May), and the end date could be later (mid-October instead of early October).

Air quality is also projected to suffer as a result of these larger, longer-lasting wildfires. Smoke from wildfires is composed of organic and black carbon particles and can impede visibility and cause respiratory problems. The U.S. Forest Service keeps a record of the amount of fuel (biomass) available across the entire United States, and another set of databases known as the Landscape Fire and Resource Management Planning Tools tracks specific types of vegetation for each square kilometer of land. Based on this information and known emission factors for combustion, the researchers predict that smoke will increase 20-100% by the 2050s, depending on the region and the type of particle.

The main innovation of the new study is its reliance on an ensemble of climate models, rather than just one or two. One of the greatest uncertainties in the science of climate change is the sensitivity of surface temperatures to rising levels of greenhouse gases.

Percentage increase in area burned, mid-21st century compared to 2000



The area burned by wildfires is expected to increase by 2050, according to research by environmental scientists at Harvard University. Credit: Xu Yue.

"Our use of a multi-model ensemble increases confidence in our results," says principal investigator Jennifer A. Logan, a recently retired Senior Research Fellow at Harvard SEAS.

The fire prediction model developed by the team performed least well in

central and southern California, where the rugged topography results in a patchwork of ecoregions, each with a different fire response to changing meteorology. The authors have been investigating the unusual factors at play in that state and expect to release their findings shortly.

For Harvard's atmospheric scientists, the ultimate goal of this project was to see how air quality could be affected by climate change, given that smoke from wildfires is a major source of particulate matter in the atmosphere.

Air quality has vastly improved over much of the United States in the past 40 years, as a result of government efforts to regulate emissions. Mickley warns that increasing wildfires may erase some of the progress.

"I think what people need to realize is that embedded in those curves showing the tiny temperature increases year after year are more extreme events that can be quite serious," she says. "It doesn't bode well."

More information: www.sciencedirect.com/science/.../S1352231013004573

Provided by Harvard University

Citation: Wildfires projected to worsen with climate change (2013, August 28) retrieved 24 April 2024 from <https://phys.org/news/2013-08-wildfires-worsen-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.