


Brown dwarf companion stars

September 2 2013



The 1.4 billion pixel camera of the Panoramic Survey Telescope & Rapid Response System (Pan-STARRS). Astronomers have used the system to study young brown dwarf stars in binary orbits around more normal young stars.
Credit: Pan-STARRS

(Phys.org) —Astronomers trying to understand how the Sun and Earth formed, and why they have their characteristic properties, have made progress on a closely related problem: the nature of the lowest mass stars, so-called "brown dwarfs." These stars have masses of less than about 8% of the Sun's mass. They are basically failed normal stars, and lack a sufficient force of gravitational contraction to heat up their

interiors to the roughly ten million kelvin temperatures needed for hydrogen burning (hydrogen burning fuels the Sun). Not surprisingly they are extremely faint and hard to detect, and as a consequence our understand of their evolution and interior properties is incomplete. Theorists predict that there could be as many brown dwarf stars as there are normal stars.

CfA [astronomer](#) Adam Kraus has joined with eight colleagues to use the Pan-STARRS1 survey (the Panoramic Survey Telescope & Rapid Response System) and an infrared survey (UKIDSS) to search in a young, 5-10 million year-old star forming region for brown dwarfs. Pan-STARRS is a wide-field imaging telescope that combines a relatively small mirror, only 1.8-meters, with a very large digital camera, about 1.4 billion pixels, to survey the sky on a continuous basis. The prototype single-mirror telescope PS1 has been operating on Mount Haleakala, Hawaii, since 2010; eventually there will be four such systems working in concert.

The astronomers were studying the very early stages of brown dwarf development, and to help obtain reliable age estimates they used these surveys to search for objects that were in a binary systems with young stars whose ages might be more tightly constrained, thus constraining the brown dwarf ages as well. The survey was able to picked out objects whose cool temperatures and reddish colors signaled that they were brown dwarfs, and the team discovered 673 binaries with candidate brown dwarf companions. Spectroscopic follow-up observations on four of them allowed the team to reach several conclusions: These companion brown dwarfs are infrequent, making up only about 0.6% of binaries, at least for those in which the stars are widely spaced (closer pairs were much more difficult to study).

More puzzling, the scientists found that the brown dwarf companions are well enough understood to conclude that they did not form at the same

time as their more normal, companion star. The puzzling result highlights the theoretical uncertainties in our understanding of brown dwarfs, while the new technique promises many more objects to begin the task of modeling. Meanwhile, the team has begun studying brown dwarfs in more normally spaced binaries, where they could be mistaken for large planets.

More information: Aller, K. et al. A Pan-STARRS + UKIDSS Search for Young, Wide Planetary-Mass Companions in Upper Scorpius, *ApJ* 773, 63, 2013.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Brown dwarf companion stars (2013, September 2) retrieved 26 June 2024 from <https://phys.org/news/2013-09-brown-dwarf-companion-stars.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--