

# Climate change: Polar bears change to diet with higher contaminant loads

September 20 2013

---



This is a harp seal with a cub. Polar bears increasingly exchange ringed seal with harp seal and hooded seal in their diet and therefore become exposed to higher concentrations of contaminants. Credit: Rune Dietz, Aarhus University

Researchers expect the climate to become warmer in the future and predict that climate change will have a significant impact on the Arctic. How will a warming Arctic affect the polar bears?

The East Greenlandic population of polar bears resides in an area, where the Arctic sea ice is expected to disappear very late. However, the decline in the ice sheet here occurs at a rate of almost 1% per year, one of the highest rates measured in the entire Arctic region.

How does this affect the prey of the polar bears - and, in turn, the polar bears' intake of contaminants? An international team of researchers set out to explore this question. The team counted researchers from the Greenland Institute of Natural Resources, Aarhus University (Denmark) and a number of Canadian institutions including: Dalhousie University, Great Lakes Institute for Environmental Research, University of Windsor, Carleton University and the National Water Research Institute.

The researchers studied the fatty acid profiles in the adipose tissue from a unique material of 310 polar bears hunted by East Greenland Inuits from the Scoresbysund area in the years from 1984 to 2011. The composition of [fatty acids](#) in the fat tissue of the polar bears namely reflects the profile of fatty acids in their diet.

The results show that the polar bears primarily feed on three species of seals: the high Arctic ringed seal and the two sub-Arctic species [harp seal](#) and hooded seal. Moreover, the results showed that the diet of the polar bears had changed over the almost 30 years during which the samples were collected. In this period, the average relative decline in the ringed seal's significance for the polar bears diet was 42%. Similarly, the intake of the sub-Arctic seals increased during the same period. Also, the researchers found that polar bears are generally in better condition now, so at a first glance the [polar bears](#) should be happy with this development.



This is a polar bear with an adult hooded seal, which make up an increasing share of their food. The hooded seal can weigh up to 300-400 kg, so it is a large prey for the approximately 500 kg heavy polar bear. Credit: Rune Dietz, Aarhus University

### **Climate change undermines improvements**

There are, however, a couple of problems that might mar the happiness, explains Professor Rune Dietz, Aarhus University:

"The problem is that the sub-Arctic seals that the polar bear has switched to, have a higher content of contaminants because they live closer to the industrialised world and are higher up in the food chain. Therefore, [climate change](#) undermines the improvements that you would otherwise have obtained owing to international regulations in the use of environmental use of persistent organic pollutants (POPs). We can see that the content of the POPs after year 2000 decreases slower in the polar bear than in, the [ringed seal](#)."

In the long term, the polar bear may very well lose access to the sub-Arctic seals as these depend on packed ice where they give birth to their cubs and are exposed to sunlight allowing them to form vital vitamin D.

**More information:** Causes and consequences of long-term change in East Greenland polar bears' diets: Investigation using quantitative fatty acid estimates and fatty acid carbon isotope patterns. McKinney, M. et al. (2013). *Global Change Biology* 19: 2360-2372. [DOI: 10.1111/gcb.12241](#)

Part 1: Three decades (1984-2010) of legacy contaminant trends in East Greenland polar bears (*Ursus maritimus*). Dietz R. et al. (2013a). *Environment International* 59:485-493. [dx.doi.org/10.1016/j.envint.2012.09.004](#)

Part 2: Three decades (1984-2010) of flame retardant trends in East Greenland polar bears (*Ursus maritimus*). Dietz R. et al. (2013b). *Environment International* 59: 494-500. [dx.doi.org/10.1016/j.envint.2012.09.008](#)

Provided by Aarhus University

Citation: Climate change: Polar bears change to diet with higher contaminant loads (2013, September 20) retrieved 22 July 2024 from <https://phys.org/news/2013-09-climate-polar-diet-higher-contaminant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.