

# Powerful jets blowing material out of galaxy: Process limits growth of central black hole and rate of star formation

September 5 2013

---

Astronomers using a worldwide network of radio telescopes have found strong evidence that a powerful jet of material propelled to nearly light speed by a galaxy's central black hole is blowing massive amounts of gas out of the galaxy. This process, they said, is limiting the growth of the black hole and the rate of star formation in the galaxy, and thus is a key to understanding how galaxies develop.

Astronomers have theorized that many [galaxies](#) should be more massive and have more stars than is actually the case. Scientists proposed two major mechanisms that would slow or halt the process of mass growth and [star formation](#)—violent stellar winds from bursts of star formation and pushback from the [jets](#) powered by the galaxy's central, supermassive black hole.

"With the finely-detailed images provided by an intercontinental combination of radio telescopes, we have been able to see massive clumps of cold gas being pushed away from the galaxy's center by the black-hole-powered jets," said Raffaella Morganti, of the Netherlands Institute for Radio Astronomy and the University of Groningen.

The scientists studied a galaxy called 4C12.50, nearly 1.5 billion light-years from Earth. They chose this galaxy because it is at a stage where the black-hole "engine" that produces the jets is just turning on. As the black hole, a concentration of mass so dense that not even light can

escape, pulls material toward it, the material forms a swirling disk surrounding the black hole. Processes in the disk tap the tremendous gravitational energy of the black hole to propel material outward from the poles of the disk.

At the ends of both jets, the researchers found clumps of hydrogen gas moving outward from the galaxy at 1,000 kilometers per second. One of the clouds has much as 16,000 times the mass of the Sun, while the other contains 140,000 times the mass of the Sun. The larger cloud, the scientists said, is roughly 160 by 190 light-years in size.

"This is the most definitive evidence yet for an interaction between the swift-moving jet of such a galaxy and a dense interstellar gas cloud," Morganti said. "We believe we are seeing in action the process by which an active, central engine can remove gas —the raw material for star formation—from a young galaxy," she added.

The scientists also said their observations indicate that the jets from the galaxy's core can stretch and deform clouds of interstellar gas to expand their "pushing" effect beyond the narrow width of the jets themselves. In addition, they reported that, at 4C12.50's stage of development, the jets may turn on and off and so periodically repeat the process of removing gas from the galaxy.

In July, another team of scientists, using the Atacama Large Millimeter/submillimeter Array (ALMA), announced they had found [gas](#) being blown from a more-nearby galaxy, called NGC 253, by an intense burst of star formation.

"Both processes are thought to be at work, often simultaneously, in young galaxies to regulate the growth of their central [black holes](#) as well as the rate at which they can form new stars," Morganti said.

The researchers published their findings in the 6 September issue of the journal *Science*.

**More information:** Radio Jets Clearing the Way Through a Galaxy: Watching Feedback in Action, *Science*, 2013.

Provided by National Radio Astronomy Observatory

Citation: Powerful jets blowing material out of galaxy: Process limits growth of central black hole and rate of star formation (2013, September 5) retrieved 23 April 2024 from <https://phys.org/news/2013-09-powerful-jets-material-galaxy-limits.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.