

Southern Ocean sampling reveals travels of marine microbes

September 18 2013



This image shows the Australia icebreaker, RSV Aurora Australis, from which samples of water in the Southern Ocean were taken at depths of up to 6 kilometers to study the marine microbes living there. Credit: UNSW

By collecting water samples up to six kilometres below the surface of the Southern Ocean, UNSW researchers have shown for the first time the impact of ocean currents on the distribution and abundance of marine

micro-organisms.

The sampling was the deepest ever undertaken from the Australian icebreaker, RSV Aurora Australis.

Microbes are so tiny they are invisible to the naked eye, but they are vital to sustaining [life on earth](#), producing most of the oxygen we breathe, soaking up carbon dioxide from the atmosphere and recycling nutrients.

"Microbes form the bulk of the biomass in oceans. All the fish, dolphins, whales, sponges and other creatures account for less than 5 per cent of the biomass," says Professor Rick Cavicchioli, of the UNSW School of Biotechnology and Biomolecular Sciences, and leader of the team.

"Microbes perform roles that nothing else can carry out. And if one critical group of [microbes](#) was destroyed, life on the planet would cease to exist."

The influence of environmental conditions on the make-up of [microbial communities](#) in different regions of the ocean has been studied, as has the role of physical barriers in preventing their dispersal.

"Collecting samples in the Southern Ocean was an enormous challenge. But it has meant we were able to carry out the first study showing how physical transport in the ocean on currents can also shape microbial communities," says Professor Cavicchioli.

The results are published in the journal *Nature Communications*.

Twenty five samples were collected across a 3000 kilometre stretch of ocean between Antarctica and the southern tip of Western Australia. Sampling depths were determined by temperature, salinity and dissolved oxygen measurements, to ensure microbes were collected from all the

distinct [water masses](#) of the Southern Ocean.

These water masses include the circumpolar deep water, which flows toward the [south pole](#) from the Indian, Pacific Ocean and Atlantic oceans; the surface water near the Antarctic coastline; and the cold, dense Antarctic bottom water, which flows north, away from the pole, at more than 4 kilometres depth.

Genetic sequencing of the microbial DNA in each sample was carried out to characterise the microbial communities in different water-masses. The research shows that communities that are connected by [ocean currents](#) are more similar to each other.

"So a microbial community could be very different to one only a few hundred metres away, but closely related to one that is thousands of kilometres away because they are connected by a current," says Professor Cavicchioli.

"Researchers need to take this into account when they are studying these important micro-organisms."

The results were tested by Dr Erik Van Sebille, of the UNSW Climate Change Research Centre, who used a computer model of the Southern Ocean to carry out a 100-year long simulation of how particles would move as a result of ocean circulation.

Dr David Wilkins, who carried out the research for his PhD under Professor Cavicchioli's supervision, is first author on the paper. The team also included UNSW's Dr Federico Lauro and Dr Stephen Rintoul, of CSIRO.

Provided by University of New South Wales

Citation: Southern Ocean sampling reveals travels of marine microbes (2013, September 18)
retrieved 24 April 2024 from

<https://phys.org/news/2013-09-southern-ocean-sampling-reveals-marine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.