


The galactic mosh pit

October 22 2013



Velocity map of the extended solar neighbourhood as seen by RAVE. Shown is a slice cut perpendicular to the plan of the Milky Way through the position of the Sun. Arrows indicate the streaming motions of the stars, the colour indicates the velocity perpendicular to the plane of the Milky Way. Credit: AIP

Astronomers have discovered that our Galaxy wobbles. An international

team of astronomers around Mary Williams from the Leibniz Institute for Astrophysics Potsdam (AIP) detected and examined this phenomenon with the RAdial Velocity Experiment (RAVE), a survey of almost half a million stars around the Sun. In addition to the regular Galactic rotation the scientists found the Milky Way moving perpendicular to the Galactic plane.

It is common knowledge that our Galaxy is permanently in motion. Being a barred [spiral galaxy](#) it rotates around the Galactic centre. It has now been discovered that our Galaxy, the Milky Way, also makes small wobbling or squishing movements. It acts like a Galactic mosh pit or a huge flag fluttering in the wind, north to south, from the Galactic plane with forces coming from multiple directions, creating a chaotic wave pattern. The source of the forces is still not understood however: possible causes include spiral arms stirring things up or ripples caused by the passage of a smaller galaxy through our own.

In this study, RAVE stars were used to examine the kinematics (velocities) of stars in a large, 3D region around the Sun - the region surveys 6500 light years above and below the Sun's position as well as inwards and outwards from the Galactic centre, reaching a quarter of the way to the centre. Using a special class of stars, red clump stars, which all have about the same brightness, mean distances to the stars could be determined. This was important as then the velocities measured with RAVE, combined with other survey data, could be used to determine the full 3D velocities (up-down, in-out and rotational). The RAVE red clump giants gave an unprecedented number of stars with which it is possible to study 3D velocities in a large region around the Sun.

The 3D movement patterns obtained showed highly complex structures. The aim was then to untangle these structures, concentrating on differences between the north and south of the Galactic plane. From these velocities it was seen that our Galaxy has a lot more going on than

previously thought. The velocities going upwards and downwards show that there is a wave-like behaviour, with [stars](#) sloshing in and out. The novel element in our approach was true 3D observation, showing how complex the velocity landscape of the Galaxy really is. Modellers now have the challenge of understanding this behaviour, be it from ripples from an eaten galaxy or the wake from [spiral arms](#). These new findings will make it possible to make 3D models of our Galaxy much more precise.

The publication "The wobbly Galaxy: kinematics north and south with RAVE red clump giants" can be found online at arxiv.org/abs/1302.2468 and was published this month in *Monthly Notices of the Royal Astronomical Society* (MNRAS).

Provided by Leibniz Institute for Astrophysics Potsdam

Citation: The galactic mosh pit (2013, October 22) retrieved 9 April 2024 from <https://phys.org/news/2013-10-galactic-mosh-pit.html>

| |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|