

# Genetics used to sort out poorly known—and hunted—whale species

October 3 2013

---



This is a rare photo of a Bryde's whale breaching in the Swatch-of-No-Ground, Bangladesh. Researchers from WCS, the American Museum of Natural History, Columbia University, and others have genetically confirmed the existence of two subspecies of Bryde's whale -- offshore and coastal forms -- in the Indian and western Pacific Oceans, important information for the management of a species hunted by Japan for scientific purposes. Credit: WCS-Bangladesh

sometimes genetically—one group of whales from another, say researchers attempting to define populations of a medium-sized and

poorly understood baleen whale that is sometimes targeted by Japan's scientific whaling program. In a new study, scientists from Wildlife Conservation Society, the American Museum of Natural History, Columbia University, NOAA, and other groups are working to define separate groups and subspecies of the Bryde's whale in the Indian and western Pacific Oceans.

By generating genetic information that allowed the team to discriminate among different Bryde's [whales](#) (almost akin to a 'bar code'), the research has confirmed the existence of two [subspecies](#), one which is larger in size and inhabits offshore waters, and one that is smaller in size and frequents more coastal marine habitats. The study now appears in the latest edition of the *Journal of Heredity*.

"Very little is known about Bryde's whales in terms of where populations are distributed, the extent of their range, or even relationships among them at the [population](#), sub-species and species levels," said Columbia University researcher Francine Kershaw, lead author of the study. "Our genetic research will help define these groups and identify populations in need of additional protection."

"The ability to delineate different populations and subspecies of Bryde's whales—particularly ones threatened by low numbers and genetic diversity—will help management authorities prevent the loss of unique and distinct genetic lineages and distinct populations," said Dr. Howard Rosenbaum, the paper's senior author and director of the Wildlife Conservation Society's Ocean Giants Program.



These are Bryde's whales lunge-feeding in the Swatch-of-No-Ground, Bangladesh. Bryde's whales are generally fish eaters, often pursuing anchovies, sardines, herring, and other schooling fish. Credit: WCS-Bangladesh

Named after Norwegian whaler and entrepreneur Johan Bryde, the Bryde's whale (pronounced BREW-dus) grows up to 50 feet in length. The species occurs in tropical, subtropical, and warm temperate waters of the Atlantic, Pacific, and Indian Oceans. In addition to scientific whaling, threats to Bryde's whale populations include ship strikes, fisheries bycatch, hydrocarbon exploration, and development in coastal waters.

The lack of knowledge on the Bryde's whale—how many species and populations there are—presents marine managers with a conservation dilemma. The species is considered "Data Deficient" by the International Union for Conservation of Nature, and the International Whaling Commission grants Japan an annual take of Bryde's whales in the

northwest Pacific Ocean through the provisions of special scientific permit. The quota for the 2012/2013 season was 34 Bryde's whales.

Working to fill in these knowledge gaps, the research team conducted an analysis of Bryde's whale populations in the Indian and Pacific Oceans by examining new genetic samples from 56 individual whales from the waters of Oman, Maldives, and Bangladesh. The team also used published data sets from Java, the coast of Japan, and the northwest Pacific in the study.

Specifically, the team examined nine diagnostic characters from the mitochondrial DNA material from each of the samples. In most instances, the researchers harmlessly collected skin samples from living whales using small biopsy darts launched from crossbows. Other samples were collected from beached whales and individuals killed in ship strikes.

The research team then examined the genetic material gathered from skin samples with a technique called polymerase chain reaction (PCR), used in this study to amplify specific regions of mitochondrial DNA (passed on through maternal lines of a population). They then carried out a statistical analysis to measure how genetically different individuals were.

The results of the genetic analysis confirmed the existence of two subspecies or types of Brydes whales, one coastal and one offshore, underlining the need to designate both subspecies as separate conservation units, with specific management needs for each type.

Further, the study revealed significant population structure for each subspecies between regions. According to the analysis, the larger offshore Bryde's whale populations in Maldives, Java, and the Northwest Pacific were genetically distinct from one another. The authors

recommend that each population be considered a separate conservation unit.

The smaller coastal form of Bryde's whale showed extremely low genetic diversity, the lowest ever measured with baleen [whale populations](#). Only a single maternal line or haplotype was detected in the 45 individual whales sampled in Bangladesh and Oman. This population, insist the authors, must be considered a conservation unit independent from coastal Bryde's whales found off Japan.

Provided by Wildlife Conservation Society

Citation: Genetics used to sort out poorly known—and hunted—whale species (2013, October 3) retrieved 23 April 2024 from

<https://phys.org/news/2013-10-genetics-poorly-knownand-huntedwhale-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.