


A link between black holes and new stars

October 23 2013



This is a composite image of NGC 1068, one of the nearest and brightest galaxies containing both a rapidly growing supermassive black hole and star formation. A new study shows that, on average, both black hole accretion and star formation grow at similar rates, perhaps related to how material flow in to the respective regions. X-ray data from the Chandra X-ray Observatory are

shown in red, optical data from the Hubble Space Telescope in green and radio data from the Very Large Array in blue. Credit: NASA and Chandra

Supermassive black holes (those with millions to billions of solar-masses) are thought to reside at the centers of most galaxies. These black holes must have undergone periods of intense accretion activity to grow to their large sizes, during which times they would be observed as active galactic nuclei and especially bright in X-rays. The masses of these black holes have been found to correlate closely with the cumulative mass of the stars that surround the galaxy in its bulge (and which are bound by gravitational forces). In a second discovery, it turns out that the activity of accretion for black holes and of star formation both peaked during a similar cosmic epoch, about three billion years ago, and then declined to the present day. These two results seem to suggest parallel evolutionary paths for the growth of the black holes and stellar mass, but the physical mechanisms that drive this apparent link, if it exists, are poorly understood. To make matters more confusing, some studies of individual objects have concluded that there is little or no correlation, or even an inverse one.

Both [black hole accretion](#) and [star formation](#) require a supply of gas, and the clues to uncovering any connections between their growth may lie in the gas fueling mechanisms that supply them. CfA astronomers Christine Jones, Bill Forman, and Andy Golding, along with a team of collaborators, reasoned that because star formation occurs over time scales of hundreds of millions of years whereas active black hole accretion can vary rapidly over millions of years, the most appropriate measure to use when comparing these two processes was a long time average.

The scientist studied 121 active galaxies studied by the Chandra X-ray

Observatory that were also observed in the infrared by the Spitzer and Herschel space telescopes. The former provided an average measure of accretion activity, while the latter, which detected the dust warmed by young stars, sampled the star formation activity. With this approach the astronomers found an almost linear relationship between the average black hole accretion rate and the star formation rate for galaxies across a wide range of luminosities. The result implies that there are indeed tight links between the two activities in galaxies.

More information: Chen, C. et al. A Correlation Between Star Formation Rate and Average Black Hole Accretion in Star-Forming Galaxies, *ApJ* 773, 3, 2013.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: A link between black holes and new stars (2013, October 23) retrieved 23 April 2024 from <https://phys.org/news/2013-10-link-black-holes-stars.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.